arXiv:1208.5139v1 [math.RT] 25 Aug 2012

MIXED SCHUR-WEYL-SERGEEV DUALITY FOR QUEER LIE SUPERALGEBRAS

JI HYE JUNG[1], SEOK-JIN KANG[2]

Abstract. We introduce a new family of superalgebras $\overrightarrow{B}_{r,s}$ for $r, s \geq 0$ such that $r + s > 0$, which we call the walled Brauer superalgebras, and prove the mixed Scur-Weyl-Sergeev duality for queer Lie superalgebras. More precisely, let $\mathfrak{q}(n)$ be the queer Lie superalgebra, $\mathbf{V} = \mathbb{C}^{n|n}$ the natural representation of $\mathfrak{q}(n)$ and $\mathbf{W}$ the dual of $\mathbf{V}$. We prove that, if $n \geq r + s$, the superalgebra $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ of the $\mathfrak{q}(n)$-action on the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. As an ingredient for the proof of our main result, we construct a new diagrammatic realization $\overrightarrow{D}_k$ of the Sergeev superalgebra Ser_k. Finally, we give a presentation of $\overrightarrow{B}_{r,s}$ in terms of generators and relations.

Introduction

The general linear group $\mathrm{GL}_n(\mathbb{C})$ of $n \times n$ invertible complex matrices acts on $V = \mathbb{C}^n$ by matrix multiplication. It is easy to see that the diagonal action of $\mathrm{GL}_n(\mathbb{C})$ is well-defined on the k-fold tensor space $V^{\otimes k}$. On the other hand, the symmetric group Σ_k acts on $V^{\otimes k}$ by place permutation. Clearly, these actions commute with each other. Moreover, Schur proved that, if $n \geq k$, then these two group actions generate the full centralizer of each other [12, 13]. This celebrated result, often referred to as the *Schur-Weyl duality*, connects the representation theories of $\mathrm{GL}_n(\mathbb{C})$ and Σ_k in a fundamental way.

There are several generalizations of the Schur-Weyl duality. For example, consider the *mixed tensor space* $V^{\otimes r} \otimes W^{\otimes s}$, where $V = \mathbb{C}^n$ is the natural representation of $\mathrm{GL}_n(\mathbb{C})$ and $W = V^*$ is its dual. The vector spaces $W^{\otimes s}$ and $V^{\otimes r} \otimes W^{\otimes s}$ inherit $\mathrm{GL}_n(\mathbb{C})$-module structures in a natural way. To describe the centralizer algebra $\mathrm{End}_{\mathrm{GL}_n(\mathbb{C})}(V^{\otimes r} \otimes W^{\otimes s})$, Koike and Turaev independently introduced the notion of *walled Brauer algebra* $B_{r,s}(n)$ [7, 16]. (See also [1].) As is the case with the pair $\mathrm{GL}_n(\mathbb{C})$ and Σ_k, the actions of $\mathrm{GL}_n(\mathbb{C})$

2000 *Mathematics Subject Classification.* 17B10, 05E10.

Key words and phrases. mixed Schur-Weyl-Sergeev duality, queer Lie superalgebras, walled Brauer superalgebras.

[1] This work was supported by NRF Grant # 2011-0027952 and by the second stage of the BK 21 Project, The Development Project of Human Resources in Mathematics, KAIST in 2011.

[2] This work was supported by NRF Grant # 2012-005700 and NRF Grant # 2011-0027952.

and $B_{r,s}(n)$ on the mixed tensor space $V^{\otimes r} \otimes W^{s}$ generate the full centralizers of each other whenever $n \geq r+s$. It is called the *mixed Schur-Weyl duality.*

On the other hand, in [15], Sergeev considered the *queer Lie superalgebra* $\mathfrak{q}(n)$ and its natural representation $\mathbf{V} = \mathbb{C}^{n|n}$. To describe the supercentralizer algebra $\operatorname{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})$, Sergeev introduced a super-extension of the symmetric group Σ_k, denoted by Ser_k, and showed that Ser_k is isomorphic to $\operatorname{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})$ whenever $n \geq k$. Moreover, he obtained a decomposition of $V^{\otimes k}$ into a direct sum of irreducible $(\mathfrak{q}(n), Ser_k)$-bimodules. The superalgebra Ser_k is called the *Sergeev superalgebra* and the duality thus obtained is called the *Schur-Weyl-Sergeev duality.* (See [8, Section 13] for more details.)

In this paper, we prove the mixed Schur-Weyl-Sergeev duality for the action of $\mathfrak{q}(n)$ on the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$, where $\mathbf{V} = \mathbb{C}^{n|n}$ is the natural representation of $\mathfrak{q}(n)$ and $\mathbf{W} = \mathbf{V}^*$ is the dual of $\mathbf{V}$. To prove our main result, we first construct a new diagrammatic realization of Ser_k, denoted by $\overrightarrow{D}_k$, which has a basis consisting of k-superdiagrams. A *k-superdiagram* is a diagram with k-vertices on its top and bottom row and the k edges connecting vertices such that each vertex on the top row is connected to exactly one vertex in the bottom row and each edge may (or may not) be marked. The multiplication on $\overrightarrow{D}_k$ is defined by *marked concatenation* of diagrams with sign. We show that the superalgebra $\overrightarrow{D}_k$ is isomorphic to Ser_k, the supercentralizer algebra $\operatorname{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})$ (Theorem 2.2).

Next, we move on to define the *walled Brauer superalgebra* $\overrightarrow{B}_{r,s}$ with a basis consisting of (r,s)-superdiagrams. An *(r,s)-superdiagram* is a diagram with $(r+s)$ vertices on the top and bottom row, the edges connecting vertices, and a vertical wall separating the r-th and $(r+1)$-th vertices in each row. Each vertex must be connected to exactly one other vertex, and each edge may (or may not) be marked. In addition, each vertical edge cannot cross the wall and each horizontal edge should cross the wall. There is a natural action of each (r,s)-superdiagram on the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$.

As a superspace, we show that $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\operatorname{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ whenever $n \geq r+s$. We give an explicit description of this linear isomorphism in Section 3. We define a multiplication on $\overrightarrow{B}_{r,s}$ by marked concatenation of diagrams with sign such that the product is zero whenever we get a loop in the middle row. Thus $\overrightarrow{B}_{r,s}$ becomes a superalgebra. It is easy to see that the even part of $\overrightarrow{B}_{r,s}$ contains the walled Brauer algebra $B_{r,s}(0)$ as a subalgebra. Our main result shows that the walled Brauer superalgebra $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\operatorname{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ whenever $n \geq r+s$ (Theorem 4.5). We also give a presentation of $\overrightarrow{B}_{r,s}$ in terms of generators and relations (Theorem 5.1).

This paper is organized as follows. In Section 1, we briefly recall the basic properties of Sergeev superalgebras. In Section 2, we construct a diagrammatic realization of Sergeev superalgebras. In Section 3, we define the superspace $\overrightarrow{B}_{r,s}$ with a basis consisting of (r, s)-superdiagrams and describe the natural action of $\overrightarrow{B}_{r,s}$ on $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. Moreover, we show that there is a linear isomorphism between $\overrightarrow{B}_{r,s}$ and $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ whenever $n \geq r + s$. In Section 4, we define a multiplication on $\overrightarrow{B}_{r,s}$ and prove our main result: the superalgebra $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ whenever $n \geq r+s$. Finally, in Section 5, we give a set of generators and defining relations for the walled Brauer superalgebra $\overrightarrow{B}_{r,s}$.

Acknowledgements. The authors would like to thank Professor Alberto Elduque at University of Zaragoza for valuable discussions on walled Brauer algebras.

1. The Sergeev superalgebras

In this paper, we will follow the notations in [8, Section 12] for the super-objects: superspaces, tensor product of superspaces, superalgebras, supermodules, superalgebra homomorphisms, supermodule homomorphisms, etc. We will also consider non-associative superalgebras.

The ground field in this paper will be $\mathbb{C}$, the field of complex numbers. We denote by $\mathbb{Z}_{\geq 0}$ the set of nonnegative integers and set $\mathbb{Z}_2 = \mathbb{Z}/2\mathbb{Z}$. For a homogeneous element v in a superspace $V = V_{\bar{0}} \oplus V_{\bar{1}}$, we write $|v| \in \mathbb{Z}_2$ for its degree.

Let Σ_k be the symmetric group of k letters which is generated by the transpositions $s_1, \ldots, s_{k-1}$.

Definition 1.1. The *Sergeev superalgebra* Ser_k is the associative superalgebra generated by $s_1, \ldots, s_{k-1}$ and $c_1, \ldots, c_k$ with the following defining relations (for admissible i, j):

$$s_i^2 = 1, \ s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \ s_i s_j = s_j s_i \qquad (|i-j| > 1), \tag{1.1}$$

$$c_i^2 = -1, \ c_i c_j = -c_j c_i \qquad (i \neq j), \tag{1.2}$$

$$s_i c_i s_i = c_{i+1}, \ s_i c_j = c_j s_i \qquad (j \neq i, i+1). \tag{1.3}$$

The generators $s_1, \ldots, s_{k-1}$ are regarded as *even* and $c_1, \ldots, c_k$ are *odd*. The subalgebra generated by $s_1, \ldots, s_{k-1}$ is isomorphic to the group algebra $\mathbb{C}\Sigma_k$ of Σ_k and the subalgebra generated by $c_1, \ldots, c_k$ is isomorphic to the *Clifford superalgebra* Cl_k. Note that Ser_k is isomorphic to the superalegbra $\mathbb{C}\Sigma_k \ltimes Cl_k$, which is $\mathbb{C}\Sigma_k \otimes Cl_k$ as a superspace with the multiplication given by

$$(\sigma \otimes c_{i_1} \cdots c_{i_\ell})(\tau \otimes c_{j_1} \cdots c_{j_m}) = \sigma\tau \otimes c_{\tau^{-1}(i_1)} \cdots c_{\tau^{-1}(i_\ell)} c_{j_1} \cdots c_{j_m}$$

for $1 \leq i_s, j_t \leq k$.

Let $\mathbf{V} = \mathbb{C}^{n|n}$ be the superspace with $\mathbf{V}_{\overline{0}} = \mathbb{C}^n$ and $\mathbf{V}_{\overline{1}} = \mathbb{C}^n$. Choose a basis $\mathbf{B}_{\overline{0}} = \{v_1, \ldots, v_n\}$ (resp. $\mathbf{B}_{\overline{1}} = \{v_{\overline{1}}, \ldots, v_{\overline{n}}\}$) of $\mathbf{V}_{\overline{0}}$ (resp. $\mathbf{V}_{\overline{1}}$). We write $|i| := |v_i| \in \mathbb{Z}_2$ for $i = 1, \ldots, n, \overline{1}, \ldots, \overline{n}$. Define an odd operator $P : \mathbf{V} \to \mathbf{V}$ by

$$v_i \longmapsto -v_{\overline{i}}, \quad v_{\overline{i}} \longmapsto v_i \quad (i = 1, \ldots, n). \tag{1.4}$$

If we abuse the notation $\overline{\overline{i}} = i$ for $i = 1, \ldots, n$, (1.4) can be written as

$$P(v_i) = (-1)^{|i|+1} v_{\overline{i}} \quad (i = 1, \ldots, n, \overline{1}, \ldots, \overline{n}).$$

Recall that the superbracket on $\mathrm{End}_{\mathbb{C}}(\mathbf{V})$ is given by

$$[f, g] = fg - (-1)^{|f||g|} gf$$

for homogeneous elements $f, g \in \mathrm{End}_{\mathbb{C}}(\mathbf{V})$. Define

$$\mathfrak{q}(n) := \{ f \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}) \mid [f, P] = 0 \}.$$

Note that it is closed under the superbracket. We call $\mathfrak{q}(n)$ the *queer Lie superalgebra*. With respect to the basis $\mathbf{B} = \mathbf{B}_{\overline{0}} \sqcup \mathbf{B}_{\overline{1}}$, we have

$$P = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$$

and the queer Lie superalgebra $\mathfrak{q}(n)$ can be expressed in the matrix form

$$\mathfrak{q}(n) = \left\{ \begin{pmatrix} A & B \\ B & A \end{pmatrix} \,\middle|\, A, B \text{ are arbitrary } n \times n \text{ complex matrices} \right\}.$$

There is a natural action of $\mathfrak{q}(n)$ on $\mathbf{V}$ by matrix multiplication, which extends to an action on the k-fold tensor product $\mathbf{V}^{\otimes k}$; i.e.,

$$\begin{aligned} & g \cdot (w_1 \otimes \cdots \otimes w_k) \\ & = \sum_{j=1}^{k} (-1)^{(|w_1| + \cdots + |w_{j-1}|)|g|} w_1 \otimes \cdots \otimes w_{j-1} \otimes g w_j \otimes w_{j+1} \otimes \cdots \otimes w_k, \end{aligned}$$

where the elements $g \in \mathfrak{q}(n)$ and $w_j \in \mathbf{V}$ $(j = 1, \cdots, k)$ are all homogeneous.

There is also a natural (right) action of the Sergeev superalgebra Ser_k on $\mathbf{V}^{\otimes k}$ [15]. Let w_j be the homogeneous element in $\mathbf{V}$. The s_j's act on $\mathbf{V}^{\otimes k}$ by graded place permutation:

$$(w_1 \otimes \cdots \otimes w_k) \cdot s_j = (-1)^{|w_j||w_{j+1}|} w_1 \otimes \cdots \otimes w_{j-1} \otimes w_{j+1} \otimes w_j \otimes w_{j+2} \otimes \cdots \otimes w_k.$$

The action of c_j's on $\mathbf{V}^{\otimes k}$ is defined as follows:

$$(w_1 \otimes \cdots \otimes w_k) \cdot c_j = (-1)^{|w_1|+\cdots+|w_{j-1}|} w_1 \otimes \cdots \otimes w_{j-1} \otimes P(w_j) \otimes w_{j+1} \otimes \cdots \otimes w_k,$$

where P is defined in (1.4). It is easy to check that the actions of s_i and c_j $(i = 1, \ldots, k-1, j = 1, \cdots, k)$ give rise to an action of the Sergeev superalgebra Ser_k on $\mathbf{V}^{\otimes k}$. Thus we obtain an algebra homomorphism

$$\Phi_k : Ser_k \longrightarrow \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes k})^{\mathrm{op}}. \tag{1.5}$$

Since Ser_k acts on $\mathbf{V}^{\otimes k}$ from the right, we consider the opposite algebra $\mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes k})^{\mathrm{op}}$.

Let $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})$ be the supercentralizer algebra of the $\mathfrak{q}(n)$-action on $\mathbf{V}^{\otimes k}$. Then we have

$$\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k}) = \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})_{\overline{0}} \oplus \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})_{\overline{1}},$$

where

$$\begin{aligned}\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})_j := \{f \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes k})_j \mid \ & f(g \cdot w) = (-1)^{|g||f|}\, g \cdot f(w) \\ & \text{for all homogeneous elements } g \in \mathfrak{q}(n) \text{ and } w \in \mathbf{V}^{\otimes k}\}.\end{aligned}$$

In [15], Sergeev proved the following fundamental theorem, the *Schur-Weyl-Sergeev duality*.

Theorem 1.2. [15, Theorem 3,4]

(a) The actions of $\mathfrak{q}(n)$ and Ser_k on $\mathbf{V}^{\otimes k}$ supercommute with each other; i.e., the image of Φ_k is in $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})$.

(b) The superalgebra homomorphism $\Phi_k : Ser_k \longrightarrow \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})^{\mathrm{op}}$ is surjective.

(c) If $n \geq k$, Φ_k is an isomorphism.

2. Diagrammatic realization of Sergeev superalgebras

In this section, we construct a new diagrammatic realization of Sergeev superalgebras. A *k-superdiagram* is defined to be a diagram consisting of k vertices on its top and bottom row together with k edges such that each vertex on the top row must be connected to exactly one vertex on the bottom row and each edge may (or may not) be *marked*. We denote the normal edge by  and the marked edge by . We number the vertices in each row of a k-superdiagram from left to right with $1, 2, \ldots, k$. In Figure 1, d is an example of a 5-superdiagram.

As usual, we can identify $\tau \in \Sigma_k$ with *its permutation diagram*, the diagram with k vertices on the top and bottom row and k normal vertical edges connecting the i-th vertex on the top row to the $\tau(i)$-th vertex (from left) on the bottom row for each $1 \leq i \leq k$.

$d =$ 

FIGURE 1. 5-superdiagram d

The k-superdiagram which has even (respectively, odd) number of marked edges is regarded as *even* (respectively, *odd*). Let $\overrightarrow{D}_k$ be the superspace with a basis consisting of k-superdiagrams. The k-superdiagrams without marked edges form a basis of $\mathbb{C}\Sigma_k$.

Now we define a multiplication on $\overrightarrow{D}_k$ in two steps.

Step 1: Marked concatenation

For $d_1, d_2 \in \overrightarrow{D}_k$, we define the *marked concatenation* $d_1 * d_2$ as follows. We first put d_1 under d_2 and identify the vertices on the bottom row of d_2 with the vertices on the top row of d_1. Next, we declare an edge in this diagram is marked if and only if the number of marked edges from d_1 and d_2 to form this edge is odd. The diagram thus obtained is the marked concatenation $d_1 * d_2$.

For example, if

$d_1 =$  , $d_2 =$  ,

then

$d_1 * d_2 =$  $=$  .

Step 2: The multiplication on $\overrightarrow{D}_k$

We first define the numbers $\rho(d_1, d_2), \ell(d_1, d_2) \in \mathbb{Z}_{\geq 0}$ as follows.

(1) Let $\rho(d_1, d_2)$ be the number of edges in $d_1 * d_2$ obtained by connecting a marked edge in d_1 and a marked edge in d_2.

(2) Consider the i-th marked edge e_i in d_1 reading the top vertices from left to right. Let a_i be the top vertex of the edge in d_2 that is connected to e_i. Then we obtain a sequence $a_1 \cdots a_p$. Let $b_1 \cdots b_q$ be the sequence obtained by reading the top vertices of the marked edges in d_2.

Consider the sequence $a_1 \cdots a_p b_1 \cdots b_q$. Suppose that the smallest entry in this sequence appears i times, and the next smallest one appears j times, etc. We replace the i occurrences of the smallest entry by $1, 2, \cdots, i$ *from left to right*, the j occurrences of the next smallest entry by $i+1, \dots, i+j$ *from left to right*, etc. We denote by $a'_1 \cdots a'_p b'_1 \cdots b'_q$ the new sequence obtained in this manner. Then we obtain a permutation $\begin{pmatrix} 1 & 2 & \cdots & p & p+1 & \cdots & p+q \\ a'_1 & a'_2 & \cdots & a'_p & b'_1 & \cdots & b'_q \end{pmatrix}$ in Σ_{p+q}. Let $\ell(d_1, d_2)$ be the length of the above permutation.

We now define the multiplication on $\overrightarrow{D}_k$ by

$$d_1 d_2 = (-1)^{\rho(d_1,d_2)+\ell(d_1,d_2)} d_1 * d_2. \tag{2.1}$$

The number $\ell(d_1, d_2)$ is called the *arranging number* for a sequence $a_1 \cdots a_p b_1 \cdots b_q$.

Example 2.1. In the above example, we have $\rho(d_1, d_2) = 1$. In (2), we obtain $a_1a_2a_3a_4 = 3125$, $b_1b_2 = 34$ and $a'_1a'_2a'_3a'_4b'_1b'_2 = 312645$. Hence $\ell(d_1, d_2) = \ell(s_2s_1s_5s_4) = 4$. It follows that

$$d_1 d_2 = (-1)^{1+4} d_1 * d_2.$$

If d_1 has i marked edges and d_2 has j marked edges, then the number of the marked edges in d_1d_2 is $i + j - 2\rho(d_1, d_2)$. It follows that $\overrightarrow{D}_k$ is a superalgebra (which may not be associative at this point). The identity element in $\overrightarrow{D}_k$ is the diagram such that each vertex on the top row is connected with the corresponding vertex on the bottom row by the normal edge.

Let d be a k-superdiagram. If we forget the marks on d, we get a permutation, say, σ. By reading the vertices at the top of marked edges from left to right, we obtain a sequence $i_1 \cdots i_m$ such that $1 \leq i_1 < \cdots < i_m \leq k$. For example, in Figure 1, we obtain

$$\sigma = s_2s_4s_1s_3 = \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 1 & 5 & 2 & 4 \end{pmatrix} \in \Sigma_5, \qquad i_1i_2i_3 = 145.$$

Recall that the Sergeev superalgebra Ser_k has a basis

$$\{\sigma c_{i_1} \cdots c_{i_m} \mid \sigma \in \Sigma_k, 1 \leq i_1 < \cdots < i_m \leq k\}. \tag{2.2}$$

Therefore we get a linear map $\phi_k : \overrightarrow{D}_k \to Ser_k$ given by

$$d \longmapsto \sigma c_{i_1} \cdots c_{i_m},$$

where σ and $i_1, \dots, i_m$ are defined above. Since $\dim \overrightarrow{D}_k = 2^k k! = \dim Ser_k$, ϕ is a linear isomorphism.

We would like to stress that, for instance, the 5-superdiagram  corresponds to an element $c_1c_2 \in Ser_5$, not c_2c_1.

We now prove the main result of this section which gives a diagrammatic realization of the Sergeev superalgebra Ser_k.

Theorem 2.2. The linear map ϕ_k preserves the multiplication:

$$\phi_k(d_1 d_2) = \phi_k(d_1)\phi_k(d_2) \quad \text{for all } d_1, d_2 \in \overrightarrow{D}_k.$$

Hence $\overrightarrow{D}_k$ is isomorphic to Ser_k as an associative superalgebra.

Proof. Since $\phi_k((\overrightarrow{D}_k)_i) \subset (Ser_k)_i$ for all $i \in \mathbb{Z}_2$, ϕ_k is an even linear map. For $d_1, d_2 \in \overrightarrow{D}_k$, let $\phi_k(d_1) = \sigma c_{i_1} \cdots c_{i_p}, \phi_k(d_2) = \tau c_{j_1} \cdots c_{j_q}$, $(\sigma, \tau \in \Sigma_k, 1 \le i_1 < \cdots < i_p \le k, 1 \le j_1 < \cdots < j_q \le k)$. Then

$$\phi_k(d_1)\phi_k(d_2) = \sigma c_{i_1} \cdots c_{i_p} \tau c_{j_1} \cdots c_{j_q} = \sigma\tau c_{\tau^{-1}(i_1)} \cdots c_{\tau^{-1}(i_p)} c_{j_1} \cdots c_{j_q}.$$

We rearrange $c_{\tau^{-1}(i_1)} \cdots c_{\tau^{-1}(i_p)} c_{j_1} \cdots c_{j_q}$ to obtain an element of the form (2.2). Since

$$c_i{}^2 = -1, c_i c_j = -c_j c_i \quad \text{for } i \neq j,$$

we have

$$c_{\tau^{-1}(i_1)} \cdots c_{\tau^{-1}(i_p)} c_{j_1} \cdots c_{j_q} = (-1)^{x+y} c_{h_1} \cdots c_{h_t},$$

where x is the number of pairs with the same entries in $\tau^{-1}(i_1) \cdots \tau^{-1}(i_p) j_1 \cdots j_q$ and y is the smallest number of transpositions that needed to rearrange $\tau^{-1}(i_1) \cdots \tau^{-1}(i_p) j_1 \cdots j_q$ in order from the smallest one to the biggest one (from left to right). We observe

$$\tau^{-1}(i_1) \cdots \tau^{-1}(i_p) j_1 \cdots j_q = a_1 \cdots a_p b_1 \cdots b_q$$

and $x = \rho(d_1, d_2)$, $y = \ell(d_1, d_2)$. Also, $\phi_k(d_1 * d_2) = \sigma\tau c_{h_1} \cdots c_{h_t}$. Therefore,

$$\begin{aligned} \phi_k(d_1)\phi_k(d_2) &= (-1)^{\rho(d_1,d_2)+\ell(d_1,d_2)} \sigma\tau c_{h_1} \cdots c_{h_t} \\ &= \phi_k(d_1 d_2). \end{aligned}$$

□

Remark 2.3. There have been a lot of works done on diagram algebras with marked edges or marked vertices (see, for example, [2, 5, 10, 11]). This work is different from those in that we deal with superalgebras.

Since $\overrightarrow{D}_k$ is isomorphic to Ser_k, it is isomorphic to $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})^{\mathrm{op}}$. Still, we will give a direct proof of this fact because it will serve as a guideline for the proof of our main theorem in Section 4.

We define a (right) action of each k-superdiagram on $\mathbf{V}^{\otimes k}$ as follows. Let $I := \{1, \ldots, n, \overline{1}, \ldots, \overline{n}\}$ and $\mathbf{i} := (i_1, \ldots, i_k) \in I^k$. Define $|i| := 0$ if $i \in \{1, \ldots, n\}$ and $|i| := 1$ if $i \in \{\overline{1}, \ldots, \overline{n}\}$. For $d \in \overrightarrow{D}_k$ and $\mathbf{i}, \mathbf{j} \in I^k$, we label the vertices at the bottom row of d with $i_1, \ldots, i_k$ (from left to right) and the vertices at the top row of d by $j_1, \ldots, j_k$ (from left to right). We denote this labeled diagram by ${}_{\mathbf{i}}d_{\mathbf{j}}$ and define the *weight* of the labeled diagram ${}_{\mathbf{i}}d_{\mathbf{j}}$ to be

$$\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) := \prod_{e} \delta_e \prod_{c} (-1)^{|c|} \prod_{\widetilde{e}_i} (-1)^{m(\widetilde{e}_i)},$$

where the notations are explained below:

(1) The first product is taken over all edges e in ${}_{\mathbf{i}}d_{\mathbf{j}}$. For each normal edge e with labels i and j at the end points, let $\delta_e = \delta_{i,j}$. We define $\delta_e = \delta_{\overline{i},j}$ for each marked edge e with label i at the bottom vertex and j at the top vertex.

(2) The second product is taken over all crossings c in ${}_{\mathbf{i}}d_{\mathbf{j}}$. We define $|c| = |e_1||e_2|$ for each crossing of two different edges e_1 and e_2, where we define $|e_i| = |i_m|$ if the edge e_i is labeled i_m at the *bottom* vertex.

(3) The last product is taken over all marked edges in ${}_{\mathbf{i}}d_{\mathbf{j}}$. Read the top vertices of the marked edges from left to right to obtain a sequence $b_1 \cdots b_q$ for $1 \leq b_1 < \cdots < b_q \leq k$ and let each marked edge be $\widetilde{e}_i$. We define $m(\widetilde{e}_i) = |j_1| + \cdots + |j_{b_i}|$.

If $\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) \neq 0$, we say that ${}_{\mathbf{i}}d_{\mathbf{j}}$ is *consistently labeled*. Since there are only vertical edges in a k-superdiagram d, the labeling $\mathbf{i} \in I^k$ determines a unique labeling $\mathbf{j} \in I^k$ such that $\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) \neq 0$.

Example 2.4. For $\mathbf{i} = (1, 2, \overline{1}, \overline{1}, \overline{2})$ and $\mathbf{j} = (1, 1, \overline{2}, \overline{2}, 1)$, the diagram ${}_{\mathbf{i}}d_{\mathbf{j}}$ of d in Figure 1 is consistently labeled and has $\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) = -1$ for $n = 2$:

${}_{\mathbf{i}}d_{\mathbf{j}} =$



.

To calculate the weight, we use

$$\prod_{c} (-1)^{|c|} = (-1)^{|1||\overline{1}| + |2||\overline{1}| + |2||\overline{2}| + |\overline{1}||\overline{2}|} = -1,$$

$$(-1)^{m(\tilde{e}_1) + m(\tilde{e}_2) + m(\tilde{e}_3)} = (-1)^{|1| + (|1| + |1| + |\overline{2}| + |\overline{2}|) + (|1| + |1| + |\overline{2}| + |\overline{2}| + |1|)} = 1.$$

Let $v_{\mathbf{i}} = v_{i_1} \otimes \cdots \otimes v_{i_k}$ for $\mathbf{i} = (i_1, \ldots, i_k) \in I^k$. Obviously, $\{v_{\mathbf{i}} \mid \mathbf{i} \in I^k\}$ forms a basis of $\mathbf{V}^{\otimes k}$. Define a linear map $\widetilde{\Phi}_k$ from $\overrightarrow{D}_k$ into $\mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes k})$ by

$$\begin{aligned} \widetilde{\Phi}_k \ : \ \overrightarrow{D}_k \ &\longrightarrow \ \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes k}) \\ d \ &\longmapsto \ \left(v_{\mathbf{i}} \mapsto \sum_{\mathbf{j} \in I^k} \mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) \ v_{\mathbf{j}} \right). \end{aligned} \tag{2.3}$$

Lemma 2.5. $\widetilde{\Phi}_k(d) = \Phi_k(\phi_k(d))$ for all $d \in \overrightarrow{D}_k$.

Proof. We observe that the action $\widetilde{\Phi}_k$ of the k-superdiagram without marked edges is the same as the action Φ_k of the corresponding permutation on $\mathbf{V}^{\otimes k}$. Assume that $d \in \overrightarrow{D}_k$ has some marked edges. If we forget the marks on d, we obtain a permutation σ. Read the top vertices of the marked edges from left to right to obtain a sequence $b_1 \cdots b_q$ for $1 \leq b_1 < \cdots < b_q \leq k$. For $\mathbf{i} \in I^k$, ${}_{\mathbf{i}}d_{\mathbf{j}}$ is consistently labeled when $\mathbf{j} = (i_{\sigma(1)}, \cdots, i_{\sigma(b_1-1)}, \overline{i_{\sigma(b_1)}}, \cdots, \overline{i_{\sigma(b_q)}}, i_{\sigma(b_q+1)}, \cdots, i_{\sigma(k)})$. Therefore, $\widetilde{\Phi}_k(d)(v_{\mathbf{i}}) = (-1)^{\prod |c|}(-1)^{m(\tilde{e}_1)+\cdots+m(\tilde{e}_q)} v_{\mathbf{j}}$, where $(-1)^{\prod |c|}$ comes from the crossing part of ${}_{\mathbf{i}}d_{\mathbf{j}}$.

By the definition, $\phi_k(d) = \sigma c_{b_1} \cdots c_{b_q}$. Therefore we have

$$\begin{aligned}
\Phi_k(\phi_k(d))\big(v_{\mathbf{i}}\big) &= \Phi_k(\sigma c_{b_1} \cdots c_{b_q})(v_{\mathbf{i}}) = \Phi_k(c_{b_q}) \cdots \Phi_k(c_{b_1}) \Phi_k(\sigma)\big(v_{\mathbf{i}}\big) \\
&= (-1)^{\prod |c|} \Phi_k(c_{b_q}) \cdots \Phi_k(c_{b_1}) \big(v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{i_{\sigma(k)}}\big) \\
&= (-1)^{\prod |c| + m(\tilde{e}_1)} \Phi_k(c_{b_q}) \cdots \Phi_k(c_{b_2}) \big(v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{\overline{i_{\sigma(b_1)}}} \otimes \cdots \otimes v_{i_{\sigma(k)}}\big) \\
&= (-1)^{\prod |c| + m(\tilde{e}_1) + \cdots + m(\tilde{e}_q)} \; v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{\overline{i_{\sigma(b_1)}}} \otimes \cdots \otimes v_{\overline{i_{\sigma(b_q)}}} \otimes \cdots \otimes v_{i_{\sigma(k)}},
\end{aligned}$$

which yields the desired result. □

Therefore, the map $\widetilde{\Phi}_k$ provides a well-defined (right) action of $\overrightarrow{D}_k$ on $\mathbf{V}^{\otimes k}$. Note that the action of the k-superdiagram without marked edges (i.e., permutation diagram) on $\mathbf{V}^{\otimes k}$ is given by graded place permutation. Moreover, we have:

Proposition 2.6.

(a) The actions of $\overrightarrow{D}_k$ and $\mathfrak{q}(n)$ on $\mathbf{V}^{\otimes k}$ supercommute with each other.
(b) The homomorphism $\widetilde{\Phi}_k : \overrightarrow{D}_k \longrightarrow \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes k})^{\mathrm{op}}$ is surjective.
(c) If $n \geq k$, $\widetilde{\Phi}_k$ is an isomorphism.

Proof. Because ϕ_k is an even homomorphism, we deduce (a) from Theorem 1.2 and Lemma 2.5. The assertions (b) and (c) follow from Theorem 1.2 and Lemma 2.5 immediately. □

3. The walled Brauer superalgebras

In this section, we will introduce a new family of superspaces $\overrightarrow{B}_{r,s}$ with a basis consisting of (r,s)-superdiagrams. We will also describe the natural (right) action of $\overrightarrow{B}_{r,s}$ on $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$ and show that it defines a linear isomorphism between $\overrightarrow{B}_{r,s}$ and $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ whenever $n \geq r+s$. Here, $\mathbf{V} = \mathbb{C}^{n|n}$ is the natural representation of $\mathfrak{q}(n)$ and $\mathbf{W} = \mathbf{V}^*$ is its dual.

To begin with, we explain the $\mathfrak{q}(n)$-supermodule structure on the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. Fix $r, s \in \mathbb{Z}_{\geq 0}$ such that $r+s > 0$. Let $I = \{1, \ldots, n, \overline{1}, \ldots, \overline{n}\}$. Recall that we fix a basis $\mathbf{B} = \{v_1, \dots, v_n, v_{\overline{1}}, \dots, v_{\overline{n}}\}$ of $\mathbf{V}$. Let us denote its dual basis by $\mathbf{B}^* = \{w_1, \dots, w_n, w_{\overline{1}}, \dots, w_{\overline{n}}\}$ of $\mathbf{W}$ such that $w_i(v_j) = \delta_{i,j}$ for $i, j \in I$. The universal enveloping algebra $U(\mathfrak{q}(n))$ is a Hopf superalgebra with the comultiplication Δ and the antipode S given by

$$\Delta(g) = g \otimes 1 + 1 \otimes g, \quad S(g) = -g \quad (g \in \mathfrak{q}(n)).$$

The dual space $\mathbf{W}$ becomes a $\mathfrak{q}(n)$-supermodule via

$$(gw)(v) := (-1)^{|g||w|} w(S(g)v)$$

for homogeneous elements $g \in \mathfrak{q}(n), w \in \mathbf{W}$ and $v \in \mathbf{V}$.

Let $E_{i,j}$ be the $n \times n$ matrix having 1 at the (i,j)-entry and 0 elsewhere. We define

$$e_{i,j} := \begin{pmatrix} E_{i,j} & 0 \\ 0 & E_{i,j} \end{pmatrix}, \quad e_{i,\overline{j}} := \begin{pmatrix} 0 & E_{i,j} \\ E_{i,j} & 0 \end{pmatrix}$$

for $1 \le i, j \le n$. We obtain the formulae

$$\begin{aligned} e_{i,j}\ v_k &= \delta_{j,k} v_i + \delta_{\overline{j},k} v_{\overline{i}}, \\ e_{i,j}\ w_k &= -(-1)^{|j||k|}(\delta_{i,k} w_j + \delta_{\overline{i},k} w_{\overline{j}}) \end{aligned} \tag{3.1}$$

for all $i \in \{1, \dots, n\}$ and $j, k \in I$. Using the comultiplication Δ, we give a $\mathfrak{q}(n)$-supermodule structure to the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. Let us denote this action by $\rho : U(\mathfrak{q}(n)) \to \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$. Then the *supercentralizer algebra* $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ of the $\mathfrak{q}(n)$-action on $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$ has a decomposition

$$\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}) := \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})_{\overline{0}} \oplus \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})_{\overline{1}},$$

where

$$\begin{aligned} \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})_j := \{ & f \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})_j \mid \\ & \rho(g) f = (-1)^{|g||f|} f \rho(g) \ \text{ for all homogeneous } g \in \mathfrak{q}(n)\}. \end{aligned}$$

Now we construct a combinatorial model for $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$. An (r,s)-*superdiagram* is a diagram with $(r+s)$ vertices on the top and bottom rows, and a vertical wall separating the r-th and $(r+1)$-th vertices (from left) in each row. Each vertex must be connected to exactly one other vertex and each edge may (or may not) be *marked*. In addition, we require that each vertical edge cannot cross the wall and each horizontal edge should cross the wall. We number the vertices in each row of an (r,s)-superdiagram from left to right with $1, 2, \dots, r+s$. The examples of $(3,2)$-superdiagrams are given in Figure 2:



FIGURE 2. $(3,2)$-superdiagrams

The (r,s)-superdiagram which has even (respectively, odd) number of marked edges is regarded as *even* (respectively, *odd*).

Let $\overrightarrow{B}_{r,s}$ be the superspace with a basis consisting of (r,s)-superdiagrams. The (r,s)-superdiagrams without marked edges form a basis of the walled Brauer algebra $B_{r,s}(\delta)$ ($\delta \in \mathbb{C}$) (See, for example, [1, 3, 4, 9, 14]).

From an $(r+s)$-superdiagram in $\overrightarrow{D}_{r+s}$, we obtain an (r,s)-superdiagram by adding the wall between the rth and $(r+1)$th vertices and interchanging the vertices on the top row with the vertices on the bottom row on the right side of the wall without disconnecting any of the edges and changing the mark of the edges. We denote this map by

$$\text{flip} : \overrightarrow{D}_{r+s} \longrightarrow \overrightarrow{B}_{r,s}. \tag{3.2}$$

For example, the $(3,2)$-superdiagrams d_1, d_2 in Figure 2 can be obtained by flipping the following 5-superdiagrams in $\overrightarrow{D}_5$, respectively:


$\text{flip}^{-1}(\mathrm{d}_1) =$
$\text{flip}^{-1}(\mathrm{d}_2) =$

It is clear that the map flip is a bijection, which implies $\dim_{\mathbb{C}} \overrightarrow{B}_{r,s} = \dim_{\mathbb{C}} \overrightarrow{D}_{r+s} = 2^{r+s}(r+s)!$.

As we did for $\overrightarrow{D}_k$ on $\mathbf{V}^{\otimes k}$ in Section 2, we will define a (right) action of each (r,s)-superdiagram in $\overrightarrow{B}_{r,s}$ on the mixed tensor space $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. In this section, we will define a linear map $\Psi_{r,s}$ from $\overrightarrow{B}_{r,s}$ to $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ and will complete the job in Section 4 by showing that $\Psi_{r,s}$ is a superalgebra homomorphism.

Given $\mathbf{i} = (i_1, \dots, i_r, i_{r+1}, \dots, i_{r+s}) \in I^{r+s}$, let $\mathbf{i}^L := (i_1, \dots, i_r)$, $\mathbf{i}^R := (i_{r+1}, \dots, i_{r+s})$ and $\mathbf{i} = \mathbf{i}^L \mathbf{i}^R$. We write $v_{\mathbf{i}} = v_{i_1} \otimes \cdots \otimes v_{i_r}$ and $w_{\mathbf{j}} = w_{j_1} \otimes \cdots \otimes w_{j_s}$ for $\mathbf{i} \in I^r, \mathbf{j} \in I^s$. Clearly, $\{v_{\mathbf{i}^L} \otimes w_{\mathbf{i}^R} \mid \mathbf{i}^L \in I^r, \mathbf{i}^R \in I^s\}$ forms a basis of $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. For $\mathbf{i}, \mathbf{j} \in I^{r+s}$ and $d \in \overrightarrow{B}_{r,s}$, we label the vertices on the bottom row of d with $i_1 \dots i_{r+s}$ (from left to right) and the vertices on the top row with d by $j_1 \dots, j_{r+s}$ (from left to right). We denote this labeled diagram by ${}_{\mathbf{i}}d_{\mathbf{j}}$. We define the *weight* of the labeled diagram ${}_{\mathbf{i}}d_{\mathbf{j}}$ to be

$$\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) := \prod_{e} \delta_e \prod_{h} (-1)^{|h|} \prod_{c} (-1)^{|c|} \prod_{e_i, \widetilde{e}_i} (-1)^{m(e_i)+m(\widetilde{e}_i)},$$

where the notations are explained below:

1. The first product is taken over all edges e in ${}_{\mathbf{i}}d_{\mathbf{j}}$. For each normal (vertical or horizontal) edge e with label i and j at the end points, let $\delta_e = \delta_{i,j}$. We define $\delta_e = \delta_{\overline{i},j}$ for each marked vertical edge e with label i at the bottom vertex and j at the top vertex, or for each marked horizontal edge e with label i at the left vertex and j at the right vertex.
2. The second product is taken over all (normal or marked) horizontal edges h on the bottom row in ${}_{\mathbf{i}}d_{\mathbf{j}}$. For each (normal or marked) horizontal edge h on the bottom row with label j at the *right* vertex of the edge, we define $|h| = |j|$.
3. The third product is taken over all the crossings c in ${}_{\mathbf{i}}d_{\mathbf{j}}$. We define $|c| = |e_1||e_2|$ for each crossing of two different edges e_1 and e_2, where we define $|e_i| = |i_m|$ if the edge e_i is a (normal or marked) vertical edge with label i_m at the *bottom* vertex, or a (normal or marked) horizontal edge with label i_m at the *right* vertex.
4. The last product is taken all marked (vertical or horizontal) edges in ${}_{\mathbf{i}}d_{\mathbf{j}}$.

   (i) Read the *left* vertices of the marked horizontal edges on the bottom row in order from left to right. Then we obtain a sequence $a_1 \cdots a_p$ for $1 \le a_1 < \cdots < a_p \le r$. Let each marked edge be e_i. We define

   $$\begin{aligned} m(e_i) &= m(e_{i-1}) + |i_{a_{i-1}+1}| + \cdots + |i_{a_i}| + 1, \\ m(e_1) &= |i_1| + \cdots + |i_{a_1}| + 1. \end{aligned}$$

   (ii) Read the *top* vertices of the marked vertical edges and the *left* vertices of the marked horizontal edges on the top row in order from left to right. Then we obtain a sequence $b_1 \cdots b_q$ for $1 \le b_1 < \cdots < b_q \le r+s$. Let each marked edge $\widetilde{e}_i$. We define $m(\widetilde{e}_i)$ as follows: (let $|j_0| = 0$)

   $$\begin{aligned} m(\widetilde{e}_i) &= |j_1| + \cdots + |j_{b_i}| && \text{if } \ b_i \le r, \\ m(\widetilde{e}_i) &= |j_1| + \cdots + |j_{(b_i)-1}| && \text{if } \ b_i > r. \end{aligned}$$

If $\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) \neq 0$, we say that ${}_{\mathbf{i}}d_{\mathbf{j}}$ is *consistently labeled.*

Example 3.1. The labeled diagram ${}_{\mathbf{i}}d_{\mathbf{j}}$ of d in Figure 3 has

$$\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}}) = \delta_{\overline{i}_1,i_3}\delta_{\overline{i}_2,j_2}\delta_{\overline{i}_4,j_3}\delta_{j_1,j_4}(-1)^{|i_3|}(-1)^{|i_2||i_3|+|i_2||j_4|+|i_4||j_4|}(-1)^{(|i_1|+1)+(|j_1|+|j_2|)+(|j_1|+|j_2|)}.$$

$d =$



FIGURE 3. $(2,2)$-superdiagram d

The above procedure provides a linear map $\Psi_{r,s}$ from $\overrightarrow{B}_{r,s}$ into $\mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$:

$$\begin{aligned} \Psi_{r,s} : \overrightarrow{B}_{r,s} &\longrightarrow \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}) \\ d &\longmapsto \left(v_{\mathbf{i}^L} \otimes w_{\mathbf{i}^R} \mapsto \sum_{\mathbf{j} \in I^{r+s}} \mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}})\ v_{\mathbf{j}^L} \otimes w_{\mathbf{j}^R} \right). \end{aligned} \tag{3.3}$$

Note that the linear map $\Psi_{r,s}$ corresponding to the (r,s)-superdiagram without marked edges is the same as the action of the walled Brauer diagram given in [3, Section 7].

Define c_i to be the following (r,s)-superdiagram:

(3.4) $c_i :=$



if $i \leq r$,

(3.5) $c_i :=$



if $i \geq r+1$.

We decompose the $(2,2)$-superdiagram d in Figure 3 into the three parts as follows:



Let d' be the $(2,2)$-superdiagram obtained from d by forgetting the marks on d. By a direct calculation, we observe

$$\Psi_{2,2}(d) = \Psi_{2,2}(c_1)\Psi_{2,2}(d')\Psi_{2,2}(c_2)\Psi_{2,2}(c_3).$$

The general case is stated in the following lemma, which can be proved by a similar calculation.

Lemma 3.2. Let d be an (r,s)-superdiagram with the sequences $a_1 \cdots a_p$ and $b_1 \cdots b_q$ which are given in (4) (i), (ii). If d' is the (r,s)-superdiagram obtained from d by forgetting the marks on d, then we have

$$\Psi_{r,s}(d) = \Psi_{r,s}(c_{a_1}) \cdots \Psi_{r,s}(c_{a_p})\Psi_{r,s}(d')\Psi_{r,s}(c_{b_1}) \cdots \Psi_{r,s}(c_{b_q}) \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}.$$

Since $\Psi_{r,s}(d')$ is even and $\Psi_{r,s}(c_j)$ is odd, we conclude that the map $\Psi_{r,s}$ is an even linear map. Moreover, we obtain

Proposition 3.3. The linear maps $\Psi_{r,s}(d)$ and $\rho(g)$ on $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$ supercommute with each other for all $d \in \overrightarrow{B}_{r,s}$ and $g \in \mathfrak{q}(n)$. That is, the image of $\Psi_{r,s}$ is contained in $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$.

Proof. By [3, Lemma 7.4], we know that the action of the (r,s)-superdiagram without the marked edges commutes with the action of $\mathfrak{q}(n)$. To prove the general case, by Lemma 3.2, it is enough to show that

$$\rho(g)\Psi_{r,s}(c_j) = (-1)^{|g|}\Psi_{r,s}(c_j)\rho(g) \tag{3.6}$$

for all homogeneous elements $g \in \mathfrak{q}(n)$.

Define the bar involution on $\mathbf{V}$ (resp. on $\mathbf{W}$) by $\overline{v_i} = v_{\bar{i}}$ (resp. $\overline{w_i} = w_{\bar{i}}$) for all $i \in I$. Using (3.1), it is easy to check that

$$\overline{e_{i,j}v_k} = e_{i,j}v_{\overline{k}}, \quad \overline{e_{i,j}w_k} = (-1)^{|j|}e_{i,j}w_{\overline{k}} \tag{3.7}$$

for all $i \in \{1, \dots, n\}$ and $j, k \in I$. Here, we use the notation $\bar{\bar{i}} = i$ for $i = 1, \dots, n$ and $\delta_{i,j} = \delta_{\bar{i},\bar{j}}$, $\delta_{\bar{i},j} = \delta_{i,\bar{j}}$ for $i, j \in I$.

Suppose $r + 1 \le j \le r + s$ and we will compare the both sides of (3.6). Set

$$v_{\mathbf{i}^L} = \mathbf{i}^L \ \text{ and } \ w_{\mathbf{i}^R} = \mathbf{i}^R \quad \text{for } \ \mathbf{i} \in I^{r+s}.$$

Then the left-hand side of (3.6) is equal to

$$\begin{aligned}
&\rho(g)\left(\Psi_{r,s}(c_j)(\mathbf{i}^L \otimes \mathbf{i}^R)\right) \\
&= \rho(g)\big((-1)^{|i_1|+\cdots+|i_{j-1}|}\, i_1 \otimes \cdots \otimes i_{j-1} \otimes \overline{i_j} \otimes i_{j+1} \otimes \cdots \otimes i_{r+s}\big) \\
&= \sum_{k<j} (-1)^{|i_1|+\cdots+|i_{j-1}|+|g|(|i_1|+\cdots+|i_{k-1}|)}\, i_1 \otimes \cdots \otimes g \cdot i_k \otimes \cdots \otimes \overline{i_j} \otimes \cdots \otimes i_{r+s} \\
&\quad + (-1)^{|i_1|+\cdots+|i_{j-1}|+|g|(|i_1|+\cdots+|i_{j-1}|)}\, i_1 \otimes \cdots \otimes g \cdot \overline{i_j} \otimes \cdots \otimes i_{r+s} \\
&\quad + \sum_{k>j} (-1)^{|i_1|+\cdots+|i_{j-1}|+|g|(|i_1|+\cdots+(|i_j|+1)+\cdots+|i_{k-1}|)}\, i_1 \otimes \cdots \otimes \overline{i_j} \otimes \cdots \otimes g \cdot i_k \otimes \cdots \otimes i_{r+s}.
\end{aligned}$$

On the other hand, the right-hand side of (3.6) is the same as

$$\begin{aligned}
&\Psi_{r,s}(c_j)\left(\rho(g)(\mathbf{i}^L \otimes \mathbf{i}^R)\right) \\
&= \Psi_{r,s}(c_j)\Big(\sum_{k=1}^{r+s} (-1)^{|g|(|i_1|+\cdots+|i_{k-1}|)}\, i_1 \otimes \cdots \otimes i_{k-1} \otimes g \cdot i_k \otimes i_{k+1} \otimes \cdots \otimes i_{r+s}\Big) \\
&= \sum_{k<j} (-1)^{|g|(|i_1|+\cdots+|i_{k-1}|)+|i_1|+\cdots+(|g|+|i_k|)+\cdots+|i_{j-1}|}\, i_1 \otimes \cdots \otimes g \cdot i_k \otimes \cdots \otimes \overline{i_j} \otimes \cdots \otimes i_{r+s} \\
&\quad + (-1)^{|g|(|i_1|+\cdots+|i_{j-1}|)+|i_1|+\cdots+|i_{j-1}|}\, i_1 \otimes \cdots \otimes \overline{g \cdot i_j} \otimes \cdots \otimes i_{r+s} \\
&\quad + \sum_{k>j} (-1)^{|g|(|i_1|+\cdots+|i_{k-1}|)+|i_1|+\cdots+|i_{j-1}|}\, i_1 \otimes \cdots \otimes \overline{i_j} \otimes \cdots \otimes g \cdot i_k \otimes \cdots \otimes i_{r+s}.
\end{aligned}$$

Now our assertion follows from (3.7).

The other case $1 \le j \le r$ can be verified in a similar manner. □

We will show that the map $\Psi_{r,s} : \overrightarrow{B}_{r,s} \to \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ is a linear isomorphism whenever $n \ge r + s$. Our argument will follow the outline given in [9, Theorem 3.4] and [3, Theorem 7.8], in which the case of $\mathfrak{gl}(m)$ and $\mathfrak{gl}(m|n)$ were treated, respectively.

Let M, N, K and L be finite dimensional $\mathfrak{q}(n)$-supermodules. We define the action of $\mathfrak{q}(n)$ on $\mathrm{Hom}_{\mathbb{C}}(M, N)$ by

$$(gf)(m) := g(f(m)) - (-1)^{|g||f|} f(gm)$$

for homogeneous elements $f \in \mathrm{Hom}_{\mathbb{C}}(M, N)$, $g \in \mathfrak{q}(n)$ and $m \in M$. Note that a linear map $f : M \to N$ is a $\mathfrak{q}(n)$-supermodule homomorphism if and only if f is annihilated by all $g \in \mathfrak{q}(n)$. We define $f^* \in \mathrm{Hom}_{\mathbb{C}}(N^*, M^*)$ by

$$(f^*(\lambda))(m) := (-1)^{|f||\lambda|}\lambda(f(m))$$

for homogeneous element $\lambda \in N^*$ and all $m \in M$. We identify

$$\mathrm{Hom}_{\mathbb{C}}(M, N) \otimes \mathrm{Hom}_{\mathbb{C}}(K, L) = \mathrm{Hom}_{\mathbb{C}}(M \otimes K, N \otimes L)$$

by

$$(f \otimes g)(m \otimes k) = (-1)^{|g||m|} f(m) \otimes g(k) \tag{3.8}$$

for $f \in \mathrm{Hom}_{\mathbb{C}}(M, N), k \in K$ and homogeneous elements $g \in \mathrm{Hom}_{\mathbb{C}}(K, L), m \in M$.

By [3, Lemma 7.4], one can deduce that the map *flip* is a $\mathfrak{q}(n)$-supermodule isomorphism:

$$\begin{array}{rccc} \textit{flip}: & \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes(r+s)}) & \longrightarrow & \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}) \\ & f \otimes g & \longmapsto & f \otimes g^*, \end{array} \tag{3.9}$$

where $f \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r})$ and $g \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes s})$.

Recall the map flip : $\overrightarrow{D}_{r+s} \longrightarrow \overrightarrow{B}_{r,s}$ given in (3.2). For a given $(r+s)$-superdiagram $d \in \overrightarrow{D}_{r+s}$, let ℓ (respectively, m) be the number of the marked vertical edges on the left-hand side (respectively, right-hand side) of the wall in flip(d). Let x (respectively, y) be the number of the marked horizontal edges on the top row (respectively, the bottom row) in flip(d).

We define

$$\begin{array}{rccc} \mathrm{Flip}: & \overrightarrow{D}_{r+s} & \longrightarrow & \overrightarrow{B}_{r,s} \\ & d & \longmapsto & (-1)^{u(d)+(\ell+x)y}\mathrm{flip}(d), \end{array}$$

where $u(d)$ is the number of crossings that are involved in the marked edges in d such that the top vertices are between the $(r+1)$th vertex and $(r+s)$th vertex.

Lemma 3.4. The following diagram is commutative:

$$\begin{tikzcd} \overrightarrow{D}_{r+s} \arrow[r, "\mathrm{Flip}"] \arrow[d, "\widetilde{\Phi}_{r+s}"] \arrow[dr, phantom, "\circlearrowleft"] & \overrightarrow{B}_{r,s} \arrow[d, "\Psi_{r,s}"] \\ \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes(r+s)}) \arrow[r, "\textit{flip}"] & \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}) \end{tikzcd} \tag{3.10}$$

Proof. For a superdiagram without the marked edges, it was proved in [3, Lemma 7.7].

For $\mathbf{i}, \mathbf{j}, \mathbf{k} \in I^r$, we define $e_{\mathbf{i},\mathbf{j}} \in \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r})$ by $e_{\mathbf{i},\mathbf{j}}(v_{\mathbf{k}}) = \delta_{\mathbf{j},\mathbf{k}} v_{\mathbf{i}}$ and for $\mathbf{i}, \mathbf{j}, \mathbf{k} \in I^s$, we define $f_{\mathbf{i},\mathbf{j}} \in \mathrm{End}_{\mathbb{C}}(\mathbf{W}^{\otimes s})$ by $f_{\mathbf{i},\mathbf{j}}(w_{\mathbf{k}}) = \delta_{\mathbf{j},\mathbf{k}} w_{\mathbf{i}}$. For $\mathbf{i} \in I^t$, let

$$|\mathbf{i}| = |i_1| + \cdots + |i_t|, \quad p(\mathbf{i}) = \sum_{1 \le a < b \le t} |i_a||i_b| \quad (\text{if } t \ge 2), \quad p(\mathbf{i}) = 0 \quad (\text{if } t = 1).$$

As in the proof of [3, Lemma 7.7], we can check that

$$w_{\mathbf{i}}(v_{\mathbf{j}}) = (-1)^{p(\mathbf{i})}\delta_{\mathbf{i},\mathbf{j}}, \quad (e_{\mathbf{i},\mathbf{j}})^* = (-1)^{(|\mathbf{i}|+|\mathbf{j}|)|\mathbf{i}|+p(\mathbf{i})+p(\mathbf{j})} f_{\mathbf{j},\mathbf{i}}. \tag{3.11}$$

From the identification (3.8), we have

$$\widetilde{\Phi}_{r+s}(d) = \sum_{\mathbf{i},\mathbf{j}\in I^{r+s}} (-1)^{(|\mathbf{i}^R|+|\mathbf{j}^R|)|\mathbf{i}^L|}\,\mathrm{wt}({}_{\mathbf{i}}d_{\mathbf{j}})e_{\mathbf{j}^L,\mathbf{i}^L}\otimes e_{\mathbf{j}^R,\mathbf{i}^R}$$

and

$$\Psi_{r,s}(\mathrm{Flip}(d)) = (-1)^{u(d)+(\ell+x)y} \sum_{\mathbf{i},\mathbf{j}\in I^{r+s}} (-1)^{(|\mathbf{i}^R|+|\mathbf{j}^R|)|\mathbf{i}^L|}\,\mathrm{wt}({}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}})e_{\mathbf{j}^L,\mathbf{i}^L}\otimes f_{\mathbf{j}^R,\mathbf{i}^R}$$

for $d \in \overrightarrow{D}_{r+s}$ and $\widetilde{d} := \mathrm{flip}(d)$. If we interchange the variables $\mathbf{i}^R$ and $\mathbf{j}^R$ and apply the *flip* map, we have

$$\mathit{flip}(\widetilde{\Phi}_{r+s}(d)) = \sum_{\mathbf{i},\mathbf{j}\in I^{r+s}} (-1)^{(|\mathbf{i}^R|+|\mathbf{j}^R|)|\mathbf{i}|+p(\mathbf{i}^R)+p(\mathbf{j}^R)}\,\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R})e_{\mathbf{j}^L,\mathbf{i}^L}\otimes f_{\mathbf{j}^R,\mathbf{i}^R}.$$

Therefore, it is enough to show that

$$(-1)^{(|\mathbf{i}^R|+|\mathbf{j}^R|)|\mathbf{i}^R|+p(\mathbf{i}^R)+p(\mathbf{j}^R)}\,\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}) = (-1)^{u(d)+(\ell+x)y}\,\mathrm{wt}({}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}}). \tag{3.12}$$

Clearly, the labeled diagram ${}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}$ is consistently labeled if and only if ${}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}}$ is consistently labeled. So we may assume $\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R})$ is not zero.

Let

$c_i :=$



$\in \overrightarrow{D}_{r+s}$ for $1 \le i \le r+s$.

If we forget the marks on d, we get a permutation σ. Note that

$$d = \sigma c_{b_1}\cdots c_{b_{\ell+x}}c_{a_1}\cdots c_{a_{y+m}},$$

where b_i, a_i are the numbers obtained by reading the top vertices of marked edges in d in order from left to right. The condition on the marked edges of $\widetilde{d}$ implies

$$1 \le b_1 < \cdots < b_{\ell+x} \le r, \ \ r+1 \le a_1 < \cdots < a_{y+m} \le r+s.$$

By the relations in $\overrightarrow{D}_{r+s}$, we obtain

$$\begin{aligned} d &= (-1)^{(\ell+x)(y+m)}\sigma c_{a_1}\cdots c_{a_{y+m}}c_{b_1}\cdots c_{b_{\ell+x}} \\ &= (-1)^{(\ell+x)(y+m)} c_{\sigma(a_1)}\cdots c_{\sigma(a_{y+m})}\sigma c_{b_1}\cdots c_{b_{\ell+x}} \\ &= (-1)^{(\ell+x)(y+m)+u(d)} c_{a'_1}\cdots c_{a'_y}c_{a''_1}\cdots c_{a''_m}\sigma c_{b_1}\cdots c_{b_{\ell+x}}, \end{aligned} \tag{3.13}$$

where $1 \leq a'_1 < \cdots < a'_y \leq r < a''_1 < \cdots < a''_m \leq r+s$. We rearrange the indices of $c_{\sigma(a_1)} \cdots c_{\sigma(a_{y+m})}$ in order from the smallest one to the biggest one (from left to right).

From Proposition 2.6 (1), we know $\widetilde{\Phi}$ preserves the multiplication on $\overrightarrow{D}_{r+s}$. Since all elements σ, c_j have only vertical edges and ${}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}$ is consistently labeled, we obtain

$$\begin{aligned}\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}) =&(-1)^{(\ell+x)(y+m)+u(d)}\,\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}c_{a'_1}{}_{\mathbf{k}^1})\cdots\mathrm{wt}({}_{\mathbf{k}^{y+m-1}}c_{a''_m}{}_{\mathbf{k}^{y+m}})\\ &\times\mathrm{wt}({}_{\mathbf{k}^{y+m}}\sigma_{\widetilde{\mathbf{k}}^1})\,\mathrm{wt}({}_{\widetilde{\mathbf{k}}^1}c_{b_1}{}_{\widetilde{\mathbf{k}}^2})\cdots\mathrm{wt}({}_{\widetilde{\mathbf{k}}^{\ell+x}}c_{b_{\ell+x}}{}_{\mathbf{j}^L\mathbf{i}^R}),\end{aligned}\tag{3.14}$$

where $\mathbf{k}^j$ (respectively, $\widetilde{\mathbf{k}}^j$) $\in I^{r+s}$ is determined by $\mathbf{k}^{j-1}$ (respectively, $\widetilde{\mathbf{k}}^{j-1}$), $\mathbf{k}^0 = {}_{\mathbf{i}^L\mathbf{j}^R}$, and $\widetilde{\mathbf{k}}^0 = \mathbf{k}^{y+m}$. Therefore, to prove (3.12), we need to calculate the right-hand side of (3.14).

(I) First, we calculate $(|\mathbf{i}^R| + |\mathbf{j}^R|)|\mathbf{i}^R| + p(\mathbf{i}^R) + p(\mathbf{j}^R)$.

Consider the partition of the set $\{r+1, \ldots, r+s\} = \{p_1 < \cdots < p_{k_1}\} \sqcup \{q_1 < \cdots < q_{k_2}\}$ such that the p_i-th vertices are the right vertices of horizontal edges on the top row in $\widetilde{d}$, and q_i-th vertices are the top vertices of vertical edges on the right-hand side of the wall in $\widetilde{d}$. Set $\mathbf{j}^{RH} := (j_{p_1}, \cdots, j_{p_{k_1}}), \mathbf{j}^{RV} := (j_{q_1}, \cdots, j_{q_{k_2}})$ for ${}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}}$. Similarly, we define $\mathbf{i}^{RH}$ and $\mathbf{i}^{RV}$ such that $\mathbf{i}^{RH}$ list the labels of the right vertices of all horizontal edges on the bottom row and $\mathbf{i}^{RV}$ list the labels of the bottom vertices of all vertical edges on the right side of the wall for ${}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}}$. Then we obtain

$$|\mathbf{j}^R| = |\mathbf{j}^{RH}| + |\mathbf{j}^{RV}|, \quad p(\mathbf{j}^R) = p(\mathbf{j}^{RH}) + |\mathbf{j}^{RH}||\mathbf{j}^{RV}| + p(\mathbf{j}^{RV}),$$

$$|\mathbf{i}^R| = |\mathbf{i}^{RH}| + |\mathbf{i}^{RV}|, \quad p(\mathbf{i}^R) = p(\mathbf{i}^{RH}) + |\mathbf{i}^{RH}||\mathbf{i}^{RV}| + p(\mathbf{i}^{RV}).$$

Since the a''_k-th vertex is the bottom vertex of the marked vertical edge in d for all $1 \leq k \leq m$, we note $\{a''_1, \ldots, a''_m\} \subseteq \{q_1, \cdots, q_{k_2}\}$ and $j_{a''_k} = \overline{i_{b''_k}}$, where b''_k-th vertex is the top vertex of the marked edge connected with a''_k-th vertex on the bottom in d. Thus we obtain the following equalities in $\mathbb{Z}_2$:

$$|\mathbf{i}^{RV}| = |\mathbf{j}^{RV}| + m,\ p(\mathbf{i}^{RV}) = p(\mathbf{j}^{RV}) + m|\mathbf{j}^{RV}| - (|j_{a''_1}| + \cdots + |j_{a''_m}|) + \binom{m}{2},$$

where $\binom{a}{b} = \dfrac{a!}{(a-b)!b!}$ and let $\binom{0}{2} = \binom{1}{2} = 0$. From the above formulae, we obtain the following equality in $\mathbb{Z}_2$:

$$\begin{aligned}&(|\mathbf{i}^R| + |\mathbf{j}^R|)|\mathbf{i}^R| + p(\mathbf{i}^R) + p(\mathbf{j}^R)\\ &= |\mathbf{i}^{RH}| + p(\mathbf{i}^{RH}) + p(\mathbf{j}^{RH}) + |\mathbf{i}^{RH}||\mathbf{j}^{RH}|\\ &\quad + m|\mathbf{j}^{RH}| + m|\mathbf{i}^{RH}| - (|j_{a''_1}| + \cdots + |j_{a''_m}|) + \binom{m}{2} + m.\end{aligned}\tag{3.15}$$

(II) Secondly, we calculate the crossing part.

An edge which does not cross the wall in $\widetilde{d}$ has the same crossings in d and $\widetilde{d}$. Observe that there are only crossing parts in $\mathrm{wt}({}_{\mathbf{k}^{y+m}}\sigma_{\widetilde{\mathbf{k}}^1})$ and

$$\mathbf{k}^{y+m} = (i_1, \dots, \overline{i_{a'_1}}, \dots, \overline{i_{a'_2}}, \dots, \overline{j_{a''_m}}, \dots, j_{r+s})$$

in (3.14). For example, consider the three crossings of the edge with labels i_2 and j_3 in the left diagram given below. We observe $\prod_c (-1)^{|c|}$ associated with the edge with labels i_2 and j_3 is $(-1)^{|i_2||i_3|+|i_2|(|i_1|+1)+|i_2||j_4|}$. In $\widetilde{d}$, we obtain $\prod_c (-1)^{|c|}$ associated with the edge with labels i_2 and j_3 is $(-1)^{|i_2||i_3|+|i_2||i_4|+|i_2||j_4|}$. Since ${}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}$ is consistently labeled, they are the same.



Since we calculate crossing parts in $\mathrm{wt}({}_{\mathbf{k}^{y+m}}\sigma_{\widetilde{\mathbf{k}}^1})$ for the left diagram above, we must multiply by $|i_1|+1$. When the edge which does not cross the wall in $\widetilde{d}$ is in the right-hand side of the wall, the situation is the same. Therefore there is no change in this part before and after flipping.

Consider two edges in d such that both edges cross the wall in $\widetilde{d}$. They cross before flipping if and only if they do not cross after flipping. For example, in the following case, the crossing of the left diagram is $(-1)^{|j_4||j_6|+|j_5||j_6|}$, and the one of the right diagram is $(-1)^{|j_4||j_5|}$, which differs by $(-1)^{p(\mathbf{j}^{RH})}$.



For the crossing of edges which become horizontal edges on the top and the bottom row in $\widetilde{d}$, respectively, it differs by $(-1)^{|\mathbf{i}^{RH}||\mathbf{j}^{RH}|}$.



The crossing part of the left diagram is $(-1)^{(|i_1|+1)|j_2|} = (-1)^{|i_2||j_2|}$ because ${}_{\mathbf{i}^L\mathbf{j}^R}d_{\mathbf{j}^L\mathbf{i}^R}$ is consistently labeled. So it differs by $(-1)^{|\mathbf{i}^{RH}||\mathbf{j}^{RH}|}$.

Similarly, we can check that for the crossing of edges which become horizontal edges on the bottom row in $\widetilde{d}$, it differs by $(-1)^{p(\mathbf{i}^{RH})}$. Therefore comparing the crossing part, we obtain the equality

$$\prod_{c}(-1)^{|c|} = \prod_{c'}(-1)^{|c'|+|\mathbf{i}^{RH}||\mathbf{j}^{RH}|+p(\mathbf{i}^{RH})+p(\mathbf{j}^{RH})} \tag{3.16}$$

for all crossings c and c' in the expression of $\mathrm{wt}({}_{\mathbf{k}^{y+m}}\sigma_{\widetilde{\mathbf{k}}^1})$ and $\mathrm{wt}({}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}})$, respectively.

(III) We calculate the part concerning with the marked edges.

In (3.13), the b_i-th vertex on the top row in d is the top vertex of a vertical marked edge $\tilde{e}$ on the left-hand side of the wall in $\widetilde{d}$ or the left vertex of a horizontal marked edge $\tilde{e}$ on the top row in $\widetilde{d}$. We observe $\mathrm{wt}({}_{\widetilde{\mathbf{k}}^1}c_{b_1}{}_{\widetilde{\mathbf{k}}^2})\cdots\mathrm{wt}({}_{\widetilde{\mathbf{k}}^{\ell+x}}c_{b_{\ell+x}}{}_{\widetilde{\mathbf{k}}^{\ell+x+1}})$ is the same as $\prod_{\tilde{e}}(-1)^{m(\tilde{e})}$ in $\mathrm{wt}({}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}})$. We also observe that the a_i'-th vertex on the bottom row is the left vertex of the horizontal marked edge e on the bottom row in $\widetilde{d}$ for all $1 \le i \le y$. Then we have $\mathrm{wt}({}_{\mathbf{i}^L\mathbf{j}^R}c_{a_1'}{}_{\mathbf{k}^1})\cdots\mathrm{wt}({}_{\mathbf{k}^{y-1}}c_{a_y'}{}_{\mathbf{k}^y})$ is the same as $\prod_e(-1)^{m(e)}$ in $\mathrm{wt}({}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}})$.

Now we consider the marked edges which are the vertical edges on the right-hand side of the wall in $\widetilde{d}$. Recall that their bottom vertices in d are $r+1 \le a_1'' < \cdots < a_m'' \le r+s$. Consider the partition of the set $\{1, \cdots, r\} = \{s_1 < \cdots < s_{k_3}\} \sqcup \{t_1 < \cdots < t_{k_4}\}$ such that the s_i-th vertices are the left vertices of the horizontal edges on the bottom in $\widetilde{d}$, and the t_i-th vertices are the bottom vertices of the vertical edges on the left-hand side of the wall in $\widetilde{d}$. Let $\mathbf{i}^{LH} := (i_{s_1}, \cdots, i_{s_{k_3}}), \mathbf{i}^{LV} := (i_{t_1}, \cdots, i_{t_{k_4}})$. Similarly, we can define $\mathbf{j}^{LH}, \mathbf{j}^{LV}$ such that $\mathbf{j}^{LH}$ list the labels of the left vertices of all horizontal edges on the top row and $\mathbf{j}^{LV}$ list the labels of the top vertices of all vertical edges on the left-hand side of the wall for ${}_{\mathbf{i}}\widetilde{d}_{\mathbf{j}}$. Then we obtain the following equalities in $\mathbb{Z}_2$:

$$|\mathbf{i}^{LV}| = |\mathbf{j}^{LV}| + \ell,\ |\mathbf{i}^{LH}| = |\mathbf{i}^{RH}| + y,\ |\mathbf{i}^{RV}| = |\mathbf{j}^{RV}| + m,\ |\mathbf{j}^{RH}| = |\mathbf{j}^{LH}| + x. \tag{3.17}$$

By a direct calculation we can check that

$$\begin{aligned} \mathrm{wt}({}_{\mathbf{k}^y}c_{a_1''}{}_{\mathbf{k}^{y+1}}) &= (-1)^{|\mathbf{i}^L|+y+|j_{r+1}|+\cdots+|j_{a_1''}|+1}, \\ \mathrm{wt}({}_{\mathbf{k}^{y+1}}c_{a_2''}{}_{\mathbf{k}^{y+2}}) &= (-1)^{|\mathbf{i}^L|+y+|j_{r+1}|+\cdots+|j_{a_1''}|+1+|j_{a_1''+1}|+\cdots+|j_{a_2''}|+1}. \end{aligned}$$

Consequently, we obtain

$$\mathrm{wt}({}_{\mathbf{k}^y}c_{a_1''}{}_{\mathbf{k}^{y+1}})\cdots\mathrm{wt}({}_{\mathbf{k}^{y+m-1}}c_{a_m''}{}_{\mathbf{k}^{y+m}}) = (-1)^{(y+|\mathbf{i}^L|)m+\sum_{k=1}^{m}\sum_{p=r+1}^{a_k''}|j_p|+\binom{m+1}{2}}.$$

From the identity (3.17), we see that the following equality holds in $\mathbb{Z}_2$:

$$(y+|\mathbf{i}^L|)m = (y+|\mathbf{i}^{LV}|+|\mathbf{i}^{LH}|)m = \ell m + m|\mathbf{j}^{LV}| + m|\mathbf{i}^{RH}|. \tag{3.18}$$

Note that $\prod_{\tilde{e}}(-1)^{m(\tilde{e})} = (-1)^{m|\mathbf{j}^L|+\sum_{k=1}^{m}\sum_{p=r+1}^{a''_k-1}|j_p|}$ for all marked vertical edges $\tilde{e}$ on the right-hand side of the wall in $\widetilde{d}$.

Combining the equalities (3.14)–(3.18), we obtain the desired result (3.12). $\square$

We now prove the main result of this section.

Theorem 3.5. The linear map $\Psi_{r,s} : \overrightarrow{B}_{r,s} \longrightarrow \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ is surjective. Moreover, when $n \geq r+s$, $\Psi_{r,s}$ is a bijection.

Proof. Since the map *flip* : $\mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes(r+s)}) \longrightarrow \mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})$ is a $\mathfrak{q}(n)$-supermodule isomorphism, our assertion follows from Proposition 2.6, Proposition 3.3, and Lemma 3.4. $\square$

4. THE MIXED SCHUR-WEYL-SERGEEV DUALITY

This section is devoted to the proof of our main result: the *mixed Schur-Weyl-Sergeev duality.* We will define a multiplication on $\overrightarrow{B}_{r,s}$ and show that $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ as an associative superalgebra whenever $n \geq r+s$.

As we did for $\overrightarrow{D}_k$, we define a multiplication on $\overrightarrow{B}_{r,s}$ in two steps.

Step 1: Marked concatenation

For $d_1, d_2 \in \overrightarrow{B}_{r,s}$, we define the *marked concatenation* $d_1 * d_2$ as follows. We first put d_1 under d_2 and identify the vertices on the bottom row of d_2 with the vertices on the top row of d_1. If there is a loop in the middle row, we define $d_1 * d_2 = 0$. If there is no loop in the middle row, we declare that an edge in this diagram is marked if and only if the number of marked edges from d_1 and d_2 to form this edge is odd. The diagram thus obtained is the marked concatenation $d_1 * d_2$.

Step 2: Multiplication on $\overrightarrow{B}_{r,s}$.

To define a multiplication, we first define the numbers $c(d_1,d_2), \ell(d_1,d_2), \rho(d_1,d_2), p(d_1,d_2) \in \mathbb{Z}_{\geq 0}$ as follows.

(1) We say that a vertex is *good* if it is the left vertex of a horizontal edge or the top vertex of a vertical edge. For a good vertex of a marked edge in d_1 that becomes a horizontal edge in the bottom row of $d_1 * d_2$, we color it with $\boxed{i}$. On the other hand, for a good vertex of a marked edge in d_1 that becomes a horizontal edge in the top row of $d_1 * d_2$, we color it with ⓘ. A good vertex of a marked edge in d_1 that becomes a vertical edge in $d_1 * d_2$ is colored with ⓘ. The numbering i

is determined by reading the good vertices from bottom to top and left to right. For the diagram d_2, we carry out the same procedure succeeding the numbering obtained from d_1. For each $\textcircled{i}$, we count the number of $\boxed{j}$'s such that $j > i$. Let $c(d_1, d_2)$ be the sum of these numbers for all $\textcircled{i}$.

(2) Let a_k be the good vertex in a new edge in $d_1 * d_2$ connected to a marked edge with color $\textcircled{i_k}$ such that $i_1 < \cdots < i_t$. Then we obtain a sequence $a_1 \cdots a_t$. Let $\ell_1(d_1, d_2)$ be the arranging number for $a_1 \cdots a_t$ defined in Section 2.

(3) Let $\rho_1(d_1, d_2)$ be the number of pairs with the same entry in $\{1, \dots, r\}$ in the sequence $a_1, \dots, a_t$.

(4) For each $\textcircled{i}$, we define the *passing number* to be the number of $\textcircled{j}$'s such that $j < i$ and $\textcircled{i}$ passes on $\textcircled{j}$ when $\textcircled{i}$ goes to a good vertex of a new edge. Let $p_1(d_1, d_2)$ be the sum of passing numbers for all $\textcircled{i}$.

(5) We carry out the same calculation of (3),(4) for the color $\boxed{i}$ to obtain the numbers $\ell_2(d_1, d_2)$ and $\rho_2(d_1, d_2)$.

(6) The *passing number* for $\boxed{i}$ is defined to be the number of $\boxed{j}$'s such that $i < j$ and $\boxed{i}$ passes on $\boxed{j}$ when $\boxed{i}$ goes to the good vertex of a new edge. Let $p_2(d_1, d_2)$ be the sum of passing numbers for all $\boxed{i}$.

(7) We define

$$\ell(d_1, d_2) := \ell_1(d_1, d_2) + \ell_2(d_1, d_2), \quad \rho(d_1, d_2) := \rho_1(d_1, d_2) + \rho_2(d_1, d_2),$$
$$\text{and} \quad p(d_1, d_2) := p_1(d_1, d_2) + p_2(d_1, d_2).$$

With these data, we define the multiplication on $\overrightarrow{B}_{r,s}$ by

$$d_1 d_2 = (-1)^{c(d_1,d_2)+\ell(d_1,d_2)+\rho(d_1,d_2)+p(d_1,d_2)} d_1 * d_2. \tag{4.1}$$

Note that if the new edge in $d_1 * d_2$ consists only of the vertical edges of d_1 and d_2, the passing number of $\textcircled{i}$ included in this new edge is always 0.

Example 4.1. If

$d_1 =$  , $d_2 =$  , then the colored diagram is given as follows:



We observe the following:

(1) The colored vertices ③, $\boxed{4}$ and $\boxed{5}$ yield $c(d_1, d_2) = 2$.
(2) The good vertices of the edges that are connected to ③, ⑥, ⑦ are $3, 3, 1$, respectively. So $\ell_1(d_1, d_2) = 2, \rho_1(d_1, d_2) = 1$.
(3) The colored vertices ③ and ⑥ yield $p_1(d_1, d_2) = 1$.
(4) The good vertices of the edges that are connected to $\boxed{1}, \boxed{2}, \boxed{4}, \boxed{5}$ are $2, 1, 1, 1$, respectively. So $\ell_2(d_1, d_2) = 3, \rho_2(d_1, d_2) = 1$.
(5) The colored vertices $\boxed{4}$ and $\boxed{5}$ yield $p_2(d_1, d_2) = 1$.

Consequently, we obtain

$$d_1 d_2 = -$$



.

From the definition of marked concatenation, we observe that $\overrightarrow{B}_{r,s}$ is a superalgebra (which may not be associative at this point). We call $\overrightarrow{B}_{r,s}$ the (r, s)-*walled Brauer superalgebra*, or simply the *walled Brauer superalgebra*. The identity element is the diagram such that each vertex on the top row is connected with the corresponding vertex on the bottom row by the normal edge.

Note that the multiplication of (r, s)-superdiagrams without marked edges is the same as the multiplication on the walled Brauer algebra $B_{r,s}(0)$. Therefore, the even part of $\overrightarrow{B}_{r,s}$ contains the walled Brauer algebra $B_{r,s}(0)$ as a subalgebra.

Now we will show that the linear map $\Psi_{r,s}$ gives a well-defined (right) action of $\overrightarrow{B}_{r,s}$ on $\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s}$. For $1 \leq p \leq r$, $r + 1 \leq q \leq r + s$, we define $e_{p,q}$ to be the following

(r, s)-superdiagram:

$e_{p,q} :=$



We use the same definition c_i given in (3.4), (3.5). A diagram in $\overrightarrow{B}_{r,s}$ with normal vertical edges only will usually be denoted by σ and will be identified with an element in $\Sigma_r \times \Sigma_s$. For simplicity, we write $\Psi_{r,s}(\sigma) = \widetilde{\sigma}$, $\Psi_{r,s}(e_{p,q}) = \widetilde{e}_{p,q}$ and $\Psi_{r,s}(c_i) = \widetilde{c}_i$.

Lemma 4.2. With the above notations, the following relations hold in $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$.

$$\begin{aligned} & \widetilde{\sigma}\widetilde{c}_j = \widetilde{c}_{\sigma(j)}\widetilde{\sigma} \text{ for } \sigma \in \Sigma_r \times \Sigma_s, \\ & \widetilde{e}_{p,q}^{\,2} = 0, \quad \widetilde{e}_{p,q}\widetilde{c}_p\widetilde{e}_{p,q} = 0, \\ & \widetilde{c}_i^{\,2} = -1 \text{ for } 1 \le i \le r, \quad \widetilde{c}_i^{\,2} = 1 \text{ for } r+1 \le i \le r+s, \\ & \widetilde{c}_i\widetilde{c}_j = -\widetilde{c}_j\widetilde{c}_i \text{ for } i \ne j, \\ & \widetilde{c}_p\widetilde{e}_{p,q} = \widetilde{c}_q\widetilde{e}_{p,q},\ \widetilde{e}_{p,q}\widetilde{c}_p = \widetilde{e}_{p,q}\widetilde{c}_q,\ \widetilde{e}_{p,q}\widetilde{c}_{p'} = \widetilde{c}_{p'}\widetilde{e}_{p,q} \text{ for different } p, q, p'. \end{aligned} \tag{4.2}$$

Proof. Each of the relations can be checked by direct calculations. For instance, we show the first relation for the case $r+1 \le j \le r+s$. Let $\mathbf{i}^L \in I^r, \mathbf{i}^R \in I^s$. We have

$$\begin{aligned} \widetilde{c}_j\widetilde{\sigma}(v_{\mathbf{i}^L} \otimes w_{\mathbf{i}^R}) =& \widetilde{c}_j((-1)^{x+y} v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{i_{\sigma(r)}} \otimes w_{i_{\sigma(r+1)}} \otimes \cdots \otimes w_{i_{\sigma(r+s)}}) \\ =& (-1)^{x+y+|\mathbf{i}^L|+|i_{\sigma(r+1)}|+\cdots+|i_{\sigma(j-1)}|} v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{i_{\sigma(r)}} \\ & \otimes w_{i_{\sigma(r+1)}} \otimes \cdots \otimes w_{\overline{i_{\sigma(j)}}} \otimes \cdots \otimes w_{i_{\sigma(r+s)}}, \end{aligned}$$

where x and y arise from crossings of the left-hand side and right-hand side of the wall in σ, respectively.

Let $\sigma(j) = k$. Then

$$\begin{aligned} \widetilde{\sigma}\widetilde{c}_k(v_{\mathbf{i}^L} \otimes w_{\mathbf{i}^R}) =& \widetilde{\sigma}((-1)^{|\mathbf{i}^L|+|i_{r+1}|+\cdots+|i_{k-1}|}\ v_{i_1} \otimes \cdots \otimes v_{i_r} \\ & \otimes w_{i_{r+1}} \otimes \cdots \otimes w_{\overline{i_k}} \otimes \cdots \otimes w_{i_{r+s}}), \\ =& (-1)^{|\mathbf{i}^L|+|i_{r+1}|+\cdots+|i_{k-1}|+x+y'} v_{i_{\sigma(1)}} \otimes \cdots \otimes v_{i_{\sigma(r)}} \\ & \otimes w_{i_{\sigma(r+1)}} \otimes \cdots \otimes w_{\overline{i_{\sigma(j)}}} \otimes \cdots \otimes w_{i_{\sigma(r+s)}}, \end{aligned}$$

where x and y' arise from crossings of the left-hand side and right-hand side of the wall in σ, respectively. One can verify that

$$(-1)^{y+|i_{\sigma(r+1)}|+\cdots+|i_{\sigma(j-1)}|} = (-1)^{y'+|i_{r+1}|+\cdots+|i_{k-1}|}.$$

There are three types of the edges of σ as shown below:



We have $y = |i_k||i_b| + |i_k||i_c| + |i_b||i_c|$ and $|i_{\sigma(r+1)}| + \cdots |i_{\sigma(j-1)}| = |i_a| + |i_c|$. Since $y' = (|i_k|+1)|i_b| + (|i_k|+1)|i_c| + |i_b||i_c|$ and $|i_{r+1}| + \cdots + |i_{k-1}| = |i_a| + |i_b|$, we obtain the desired result. □

The following proposition is one of the key ingredients in proving our main theorem.

Proposition 4.3. The linear map $\Psi_{r,s}$ defines a superalgebra homomorphism between $\overrightarrow{B}_{r,s}$ and $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$. That is, for all $d_1, d_2 \in \overrightarrow{B}_{r,s}$, we have

$$\Psi_{r,s}(d_1 d_2) = \Psi_{r,s}(d_1)\Psi_{r,s}(d_2) \quad \text{in} \quad \mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}.$$

Proof. By [3, Lemma 7.4], the map $\Psi_{r,s}$ preserves the multiplication for (r,s)-superdiagrams without marked edges.

First, we assume that there is a loop in the middle row of $d_1 * d_2$. If the number of marked edges to form a loop is odd, there is no $\mathbf{j}$ such that ${}_{\mathbf{i}}d_{1\mathbf{j}}$, ${}_{\mathbf{j}}d_{2\mathbf{k}}$ are all consistently labeled. For any starting point of the loop, we label j and $\overline{j}$ simultaneously since there is an odd number of the horizontal edges. Therefore, $\sum_{\mathbf{j} \in I^{r+s}} \mathrm{wt}({}_{\mathbf{i}}d_{1\mathbf{j}})\,\mathrm{wt}({}_{\mathbf{j}}d_{2\mathbf{k}}) = 0$ for all $\mathbf{k} \in I^{r+s}$, and hence $\Psi_{r,s}(d_1)\Psi_{r,s}(d_2) = 0$.

Suppose the number of marked edges to form a loop is even. Simply, we write $\widetilde{x} = \Psi_{r,s}(x)$ for $x \in \overrightarrow{B}_{r,s}$. Let $\widetilde{c}_A := \widetilde{c}_{a_1} \cdots \widetilde{c}_{a_m}$ for a non-empty subset $A = \{a_1 < \cdots < a_m\} \subset \{1, \cdots, r+s\}$, and $\widetilde{c}_A := 1$ for an empty set A. Using the decomposition in Lemma 3.2, we obtain

$$\widetilde{d}_1 = \widetilde{c}_{a_1} \cdots \widetilde{c}_{a_p} \widetilde{d}'_1 \widetilde{c}_{b_1} \cdots \widetilde{c}_{b_q}, \quad \widetilde{d}_2 = \widetilde{c}_{a'_1} \cdots \widetilde{c}_{a'_t} \widetilde{d}'_2 \widetilde{c}_{b'_1} \cdots \widetilde{c}_{b'_u}.$$

Since the proof for the general case is too messy, we will work with the following example to explain the main idea of proof. Let us draw only a loop in the middle row in $d_1 * d_2$

and let the vertices in the loop be $x_1, x_2, x_3, y_1, y_2, y_3$ from left to right.

$d_1 * d_2 = \quad \cdots$



$\cdots$

Observe that

$$d_1' = \sigma e_{x_1,y_1} e_{x_2,y_3} e_{x_3,y_2}, \quad d_2' = e_{x_1,y_3} e_{x_2,y_2} e_{x_3,y_1} \tau$$

for some (r,s)-superdiagrams σ, τ without the marked edges. Since $\Psi_{r,s}$ preserves the multiplication for (r,s)-superdiagrams without the marked edges, combined with Lemma 4.2, we obtain

$$\widetilde{d_1} = \pm \widetilde{c}_{A_1} \widetilde{\sigma} \widetilde{c}_{B_1} \widetilde{e}_{x_1,y_1} \widetilde{e}_{x_2,y_3} \widetilde{e}_{x_3,y_2} \widetilde{c}_{x_2}, \quad \widetilde{d_2} = \pm \widetilde{c}_{x_3} \tilde{e}_{x_1,y_3} \tilde{e}_{x_2,y_2} \tilde{e}_{x_3,y_1} \widetilde{c}_{A_2} \widetilde{\tau} \widetilde{c}_{B_2},$$

where A_i, B_i are the set of indices of the marked edges. Since $\widetilde{c}_i \widetilde{c}_j = -\widetilde{c}_j \widetilde{c}_i$ for $i \neq j$, we need to take the sign into account.

By the associativity of $\mathrm{End}_{\mathbb{C}}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ and the relations in Lemma 4.2, we know $\widetilde{c}_{x_3}$ can be transformed into $\widetilde{c}_{x_2}$ in $\widetilde{e}_{x_1,y_1} \widetilde{e}_{x_2,y_3} \widetilde{e}_{x_3,y_2} \widetilde{c}_{x_2} \widetilde{c}_{x_3} \tilde{e}_{x_1,y_3} \tilde{e}_{x_2,y_2} \tilde{e}_{x_3,y_1}$. In general case, since there are even number of the marked edges in a loop, the $\widetilde{c}_i$'s will be canceled out, too. Notice that

$$\widetilde{e}_{x_1,y_1} \widetilde{e}_{x_2,y_3} \widetilde{e}_{x_3,y_2} \tilde{e}_{x_1,y_3} \tilde{e}_{x_2,y_2} \tilde{e}_{x_3,y_1} = \Psi_{r,s}(e_{x_1,y_1} e_{x_2,y_3} e_{x_1,y_3} e_{x_2,y_2} e_{x_3,y_1}) = 0.$$

Since $\widetilde{d_1} \widetilde{d_2} = \pm \cdots \widetilde{e}_{x_1,y_1} \widetilde{e}_{x_2,y_3} \widetilde{e}_{x_3,y_2} \widetilde{c}_{x_2} \widetilde{c}_{x_3} \tilde{e}_{x_1,y_3} \tilde{e}_{x_2,y_2} \tilde{e}_{x_3,y_1} \cdots$, we obtain the desired result.

The general case can be handled in this manner.

Next, let us consider the case $d_1 d_2 \neq 0$. For an (r,s)-superdiagram d without marked edges, we can decompose d as follows.

(1) For d in $\overrightarrow{B}_{r,s}$, read the left vertex of each horizontal edges on the bottom row from left to right to obtain a sequence $p_1 \cdots p_a$. We denote by q_a the right vertex of the horizontal edge connected with p_a-th vertex.

(2) Read the left vertex of each horizontal edge on the top row from left to right to obtain a sequence $p_1' \cdots p_a'$. Let q_a' be the right vertex of the edge connected with p_a'-th vertex.

(3) Let σ be the (r,s)-superdiagram such that p_i'-th vertex (respectively, q_i'-th vertex) on the top row is connected with p_i-th vertex (respectively, q_i-th vertex) on the bottom row by a normal edge. The other edges of σ are all normal and have the same connection

as d. Therefore, $\sigma \in \Sigma_r \times \Sigma_s$ and $\sigma^{-1}(p_i) = p'_i$, $\sigma^{-1}(q_i) = q'_i$ for all i, which implies $d = e_{p_1,q_1} \cdots e_{p_a,q_a}\sigma$.

For a general $d \in \overrightarrow{B}_{r,s}$, let P (respectively, Q) be the set of good vertices of marked horizontal edges on the bottom row (respectively, marked vertical edges or marked horizontal edges on the top row) of d. Let $c_A := (((c_{a_1}c_{a_2})\cdots)c_{a_m})$ for a non-empty set $A = \{a_1 < \cdots < a_m\} \subset \{1, \cdots, r+s\}$, and $c_A := 1$ for an empty set A, where c_i is the (r,s)-superdiagram defined in (3.4), (3.5). Then we can decompose d in the following form:

$$d = ((((c_P e_{p_1,q_1})\cdots e_{p_a,q_a})\sigma)c_Q) \tag{4.3}$$

satisfying the conditions

(i) $1 \le p_1 < \cdots < p_a \le r$, and $r+1 \le q_i \le r+s$ are all distinct,

(ii) $\sigma \in \Sigma_r \times \Sigma_s$ and $\sigma^{-1}(p_1) < \cdots < \sigma^{-1}(p_a)$,

(iii) $P \subset \{p_1, \cdots, p_a\}$, $Q \subset \{1, \cdots, r+s\} \setminus \{\sigma^{-1}(q_1), \cdots, \sigma^{-1}(q_a)\}$.

Note that when we calculate the multiplication in the right-hand side of (4.3), the exponent of -1 is always 0. By Lemma 3.2 and [3, Lemma 7.4], we obtain the decomposition

$$\widetilde{d}_1 = \widetilde{c}_P \widetilde{e}_{p_1,q_1} \cdots \widetilde{e}_{p_a,q_a} \widetilde{\sigma} \widetilde{c}_Q, \quad \widetilde{d}_2 = \widetilde{c}_T \widetilde{e}_{t_1,u_1} \cdots \widetilde{e}_{t_x,u_x} \widetilde{\tau} \widetilde{c}_U.$$

For simplicity, let $\widetilde{d}'_1 = \widetilde{e}_{p_1,q_1} \cdots \widetilde{e}_{p_a,q_a}\widetilde{\sigma}$, $\widetilde{d}'_2 = \widetilde{e}_{t_1,u_1} \cdots \widetilde{e}_{t_x,u_x}\widetilde{\tau}$.

Note that we color all the elements of P, Q, T, U with $\boxed{i}$ or ⓘ. Also we color the elements of P (respectively, U) with the square (respectively, circle). Let A_1 (respectively, A_2) be the set of elements in Q which are colored with the square (respectively circle). Similarly, let B_1 (respectively, B_2) be the set of elements in T which are colored with the square (respectively, circle). Then we have

$$\widetilde{d}_1 \widetilde{d}_2 = (-1)^{c(d_1,d_2)} \widetilde{c}_P \widetilde{d}'_1 \widetilde{c}_{A_1} \widetilde{c}_{B_1} \widetilde{c}_{A_2} \widetilde{c}_{B_2} \widetilde{d}'_2 \widetilde{c}_U.$$

We shift the element $\widetilde{c}_i$ $(i \in A_2, B_2, U)$ using the relations in Lemma 4.2. Then we have

$$\widetilde{d}_1 \widetilde{d}_2 = (-1)^{c(d_1,d_2)+p_1(d_1,d_2)} \widetilde{c}_P \widetilde{d}'_1 \widetilde{c}_{A_1} \widetilde{c}_{B_1} \widetilde{d}'_2 \widetilde{c}_{a_1} \cdots \widetilde{c}_{a_t},$$

where a_i's are the good vertices of new edges in $d_1 * d_2$ and $p_1(d_1,d_2)$ is the sum of passing numbers for all ⓘ. The exponent $p_1(d_1,d_2)$ arises from the following formula:

$$\widetilde{e}_{p,q} \widetilde{c}_{b_1} \cdots \widetilde{c}_{b_m} \widetilde{c}_q = (-1)^x \widetilde{e}_{p,q} \widetilde{c}_{b_1} \cdots \widetilde{c}_{b_m} \widetilde{c}_p,$$

where all $b_i \neq q$ and x is the number of b_i's such that $b_i = p$. For example, if

$d_1 =$  and $d_2 =$  ,

the labeled diagram is as follows:



.

By Lemma 4.2, we obtain

$$\begin{aligned}\widetilde{d}_1\widetilde{d}_2 &= (\tilde{e}_{2,3}\widetilde{c}_2)(\widetilde{c}_1\tilde{e}_{1,3}\widetilde{s}_1) = \tilde{e}_{2,3}\widetilde{c}_2(\widetilde{c}_1\tilde{e}_{1,3})\widetilde{s}_1 = \tilde{e}_{2,3}\widetilde{c}_2(\widetilde{c}_3\tilde{e}_{1,3})\widetilde{s}_1 = (\tilde{e}_{2,3}\widetilde{c}_2\widetilde{c}_3)\tilde{e}_{1,3}\widetilde{s}_1\\ &= (-1)\ (\tilde{e}_{2,3}\widetilde{c}_2\widetilde{c}_2)\tilde{e}_{1,3}\widetilde{s}_1 = (-1)\ \tilde{e}_{2,3}\widetilde{c}_2(\widetilde{c}_2\tilde{e}_{1,3}\widetilde{s}_1) = (-1)\ \tilde{e}_{2,3}\widetilde{c}_2(\tilde{e}_{1,3}\widetilde{s}_1\widetilde{c}_1)\\ &= (-1)\ \tilde{e}_{2,3}(\widetilde{c}_2\tilde{e}_{1,3}\widetilde{s}_1)\widetilde{c}_1 = (-1)\ \tilde{e}_{2,3}(\tilde{e}_{1,3}\widetilde{s}_1\widetilde{c}_1)\widetilde{c}_1 = (-1)\ \tilde{e}_{2,3}\tilde{e}_{1,3}\widetilde{s}_1\widetilde{c}_1\widetilde{c}_1.\end{aligned}$$

Moreover, we have

$$\widetilde{c}_{a_1}\cdots\widetilde{c}_{a_t} = (-1)^{\ell_1(d_1,d_2)+\rho_1(d_1,d_2)}\widetilde{c}_{h'_1}\cdots\widetilde{c}_{h'_z},$$

where $h'_1 < h'_2 < \cdots < h'_z$. It follows that

$$\widetilde{d}_1\widetilde{d}_2 = (-1)^{c(d_1,d_2)+p_1(d_1,d_2)+\ell_1(d_1,d_2)+\rho_1(d_1,d_2)}\widetilde{c}_P\widetilde{d}'_1\widetilde{c}_{A_1}\widetilde{c}_{B_1}\widetilde{d}'_2\widetilde{c}_{h'_1}\cdots\widetilde{c}_{h'_z}.$$

Similarly, by shifting $\widetilde{c}_i$'s for $i \in P, A_1, B_1$, we have

$$\widetilde{d}_1\widetilde{d}_2 = (-1)^{c(d_1,d_2)+p(d_1,d_2)+\ell(d_1,d_2)+\rho(d_1,d_2)}\widetilde{c}_{h_1}\cdots\widetilde{c}_{h_y}\widetilde{d}'_1\widetilde{d}'_2\widetilde{c}_{h'_1}\cdots\widetilde{c}_{h'_z},$$

where $h_1 < h_2 < \cdots < h_y$. Note that $\Psi_{r,s}(d_1 * d_2) = \widetilde{c}_{h_1}\cdots\widetilde{c}_{h_y}\widetilde{d}'_1\widetilde{d}'_2\widetilde{c}_{h'_1}\cdots\widetilde{c}_{h'_z}$. Hence we obtain

$$\Psi_{r,s}(d_1)\Psi_{r,s}(d_2) = \Psi_{r,s}(d_1d_2)$$

as desired. □

Example 4.4. For $(3,3)$-superdiagrams d_1, d_2 in Example 4.1, we can decompose

$$d_1 = c_2e_{2,5}s_4c_1c_2c_6,\ d_2 = c_1c_3e_{1,6}e_{3,4}s_2c_1.$$

The decomposition of d_1 is shown below:



Now we are ready to state and prove our main theorem.

Theorem 4.5.
(a) The walled Brauer superalgebra $\overrightarrow{B}_{r,s}$ is an associative superalgebra for all $r, s \geq 0$.

(b) When $n \geq r + s$, the walled Brauer superalgebra $\overrightarrow{B}_{r,s}$ is isomorphic to the supercentralizer algebra $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ as an associative superalgebra.

Proof. (a) When $n \geq r + s$, since $\mathrm{End}_{\mathfrak{q}(n)}(\mathbf{V}^{\otimes r} \otimes \mathbf{W}^{\otimes s})^{\mathrm{op}}$ is an associative superalgebra, Theorem 3.5 and Theorem 4.3 show that $\overrightarrow{B}_{r,s}$ is an associative superalgebra. Note that the associativity condition $(d_1 d_2)d_3 = d_1(d_2 d_3)$ does not depend on n for all (r, s)-superdiagrams $d_1, d_2, d_3 \in \overrightarrow{B}_{r,s}$. Therefore, if $\overrightarrow{B}_{r,s}$ is associative for large enough n, then $\overrightarrow{B}_{r,s}$ is associative for all $r, s \geq 0$.

Now the assertion (b) is obvious. □

5. Presentation of walled Brauer superalgebras

Another main result of this paper is a presentation of $\overrightarrow{B}_{r,s}$. Assume that $1 \leq i \leq r - 1$, $r + 1 \leq j \leq r + s - 1$, $1 \leq k \leq r$, $r + 1 \leq l \leq r + s$. Let us consider the following diagrams in $\overrightarrow{B}_{r,s}$:



$s_j :=$  ,

$e_{r,r+1} :=$  ,

$c_k :=$  ,

$c_l :=$  .

Theorem 5.1. The walled Brauer superalgebra $\overrightarrow{B}_{r,s}$ is the associative superalgebra generated by (even generators) $1, s_1, \ldots, s_{r-1}, s_{r+1}, \ldots, s_{r+s-1}, e_{r,r+1}$ and (odd generators) $c_1, \ldots, c_{r+s}$ with the following defining relations (for admissible i, j) :

$$s_i^2 = 1, \quad s_i s_{i+1} s_i = s_{i+1} s_i s_{i+1}, \quad s_i s_j = s_j s_i \quad (|i-j| > 1), \tag{5.1}$$

$$e_{r,r+1}^2 = 0, \quad e_{r,r+1} s_j = s_j e_{r,r+1} \quad (j \neq r-1, r+1), \tag{5.2}$$

$$e_{r,r+1} = e_{r,r+1} s_{r-1} e_{r,r+1} = e_{r,r+1} s_{r+1} e_{r,r+1}, \tag{5.3}$$

$$s_{r-1} s_{r+1} e_{r,r+1} s_{r+1} s_{r-1} e_{r,r+1} = e_{r,r+1} s_{r-1} s_{r+1} e_{r,r+1} s_{r+1} s_{r-1}, \tag{5.4}$$

$$c_i^2 = -1 \quad (1 \leq i \leq r), \quad c_i^2 = 1 \quad (r+1 \leq i \leq r+s), \quad c_i c_j = -c_j c_i \quad (i \neq j), \tag{5.5}$$

$$s_i c_i s_i = c_{i+1}, \quad s_i c_j = c_j s_i \quad (j \neq i, i+1), \tag{5.6}$$

$$c_r e_{r,r+1} = c_{r+1} e_{r,r+1}, \quad e_{r,r+1} c_r = e_{r,r+1} c_{r+1}, \tag{5.7}$$

$$e_{r,r+1} c_r e_{r,r+1} = 0, \quad e_{r,r+1} c_j = c_j e_{r,r+1} \quad (j \neq r, r+1). \tag{5.8}$$

Proof. We observe that the subalgebra of $\overrightarrow{B}_{r,s}$ generated by s_i's ($i = 1, \ldots, r-1, r+1, \ldots, r+s-1$) is isomorphic to the group algebra of $\Sigma_r \times \Sigma_s$. Also we have $\sigma e_{r,r+1} \sigma^{-1} = e_{\sigma(r),\sigma(r+1)}$ for $\sigma \in \Sigma_r \times \Sigma_s$. Since every element $d \in \overrightarrow{B}_{r,s}$ can be decomposed into the form (4.3), the elements $s_1, \ldots, s_{r-1}, s_{r+1}, \ldots, s_{r+s-1}, e_{r,r+1}$ and $c_1, \ldots, c_{r+s}$ generate the associative superalgebra $\overrightarrow{B}_{r,s}$. By direct calculations, one can verify that these elements satisfy the above relations.

Now take the associative superalgebra F over $\mathbb{C}$ generated by s_i ($i = 1, \ldots, r-1, r+1, \ldots, r+s-1$), $e_{r,r+1}$ and c_j ($1 \leq j \leq r+s$) with the above defining relations. Then $\overrightarrow{B}_{r,s}$ is a quotient of F. In particular, $\dim_{\mathbb{C}}(\overrightarrow{B}_{r,s}) \leq \dim_{\mathbb{C}} F$. Therefore, to prove our theorem, it is enough to show that $\dim_{\mathbb{C}} F \leq \dim_{\mathbb{C}}(\overrightarrow{B}_{r,s})$.

Let Σ be the subalgebra of F generated by s_i's and let $\widetilde{s}_i$ be the transposition in $\Sigma_r \times \Sigma_s$ corresponding to $i = 1, \ldots, r-1, r+1, \ldots, r+s-1$. We define an algebra homomorphism $\psi : \mathbb{C}(\Sigma_r \times \Sigma_s) \longrightarrow \Sigma$ by $\widetilde{s}_i \mapsto s_i$. It is a well-defined surjective homomorphism. So Σ is a quotient of $\mathbb{C}(\Sigma_r \times \Sigma_s)$. Let $e_{p,q} := \sigma e_{r,r+1} \sigma^{-1}$, where $\sigma = s_{q-1} \cdots s_{r+1} s_p \cdots s_{r-1}$ for $1 \leq p \leq r-1$, $r+2 \leq q \leq r+s$. Set $c_A := c_{a_1} \cdots c_{a_m}$ for a non-empty set $A = \{a_1 < \cdots < a_m\} \subset \{1, \cdots, r+s\}$ and $c_A := 1$ for an empty set A. We define

$$\begin{aligned} X := \{c_P e_{p_1,q_1} \cdots e_{p_a,q_a} c_Q \sigma \mid\ & \text{(i) } 1 \leq p_1 < \cdots < p_a \leq r, \\ & \text{(ii) } r+1 \leq q_i \leq r+s \text{ and } q_i \text{ are all distinct}, \\ & \text{(iii) } \sigma \in \Sigma, \text{ and } {}^{\exists}\widetilde{\sigma} \text{ such that } \psi(\widetilde{\sigma}) = \sigma, \text{ and } \widetilde{\sigma}^{-1}(p_1) < \cdots < \widetilde{\sigma}^{-1}(p_a), \\ & \text{(iv) } P \subset \{p_1, \ldots, p_a\},\ Q \subset \{1, \ldots, r+s\} \setminus \{q_1, \ldots, q_a\}\}. \end{aligned}$$

We will prove our assertion in two steps:

Step 1: Every word in the generators of F belongs to the linear span of elements of X.

Step 2: The number of elements in X is less than or equal to the dimension of $\overrightarrow{B}_{r,s}$.

Proof of Step 1:

For $1 \leq i < j \leq r$ or $r+1 \leq i < j \leq r+s$, we define

$$(i\ j) := s_i s_{i+1} \cdots s_{j-2} s_{j-1} s_{j-2} \cdots s_{i+1} s_i \in \Sigma.$$

For $1 \leq j < i \leq r$ or $r+1 \leq j < i \leq r+s$, define $(i\ j) := (j\ i)^{-1} \in \Sigma$. From the relations (5.1)–(5.8), we obtain the following relations:

$$\begin{aligned} &(1) \quad (p'\ q') e_{p,q} = e_{p,q} (p'\ q'), \\ &(2) \quad (p\ p') e_{p,q} = e_{p',q} (p\ p'),\ \ (q\ q') e_{p,q} = e_{p,q'} (q\ q'), \\ &(3) \quad e_{p,q} e_{p',q'} = e_{p',q'} e_{p,q}, \\ &(4) \quad e_{p,q} e_{p,q'} = e_{p,q} (q\ q'),\ \ e_{p,q} e_{p',q} = e_{p,q} (p\ p'), \\ &(5) \quad e_{p,q} e_{p',q'} (p\ p')(q\ q') = e_{p,q} e_{p',q'}, \\ &(6) \quad c_q e_{p,q} = c_p e_{p,q},\ \ e_{p,q} c_q = e_{p,q} c_p, \\ &(7) \quad c_{p'} e_{p,q} = e_{p,q} c_{p'}, \\ &(8) \quad e_{p,q}^2 = 0, \quad e_{p,q} c_p e_{p,q} = 0. \end{aligned}$$

Here, all p, q, p', q' are admissible and distinct.

Suppose that the relations (1)–(8) are satisfied. For a given word w in the generators of F, using the relations (5.6), we can interchange the place of c_i with the place of s_j. From the relation (5.1) and the definition of $e_{p,q}$, we may write w in terms of the elements $e_{p,q}$. From the relations (1) and (2), we may move $e_{p,q}$ in the left-hand side of all s_i's. We would like to move c_i in the left-hand side of all $e_{p,q}$'s or in the right-hand side of all $e_{p,q}$'s.

From the relation (7), we move each c_j in the left-most side of w, except for c_p's which are in the right-hand side of all $e_{p,q}$'s. By the relations (5.5) and (6), we have

$$w = \pm c_X e_{a_1,b_1} \cdots e_{a_\ell,b_\ell} e_{i_1,i'_1} c_{i_1} e_{a'_1,b'_1} \cdots e_{a'_x,b'_x} e_{i_2,i'_2} c_{i_2} \cdots$$

for some set X. Using the relations (1)–(4) and (8), we rearrange $e_{a_1,b_1} \cdots e_{a_\ell,b_\ell} e_{i_1,i'_1}$ to obtain $e_{h_1,g_1} \cdots e_{h_m,g_m} s_{x_1} \cdots s_{x_u}$ or 0, where h_j, g_j are all distinct. From the relation (5.6), we obtain $s_{x_1} \cdots s_{x_m} c_{i_1} = c_{k_1} s_{x_1} \cdots s_{x_m}$ for some k_1. If all h_j is not k_1, then we can shift c_{k_1} to the left-hand side of $e_{h_1,g_1} \cdots e_{h_m,g_m}$. If $h_j = k_1$ for some j, e_{h_p,g_p} $(p \neq j)$ can be shifted to the right-hand side of c_{k_1}. In this way, we can obtain $w = 0$ or

$$w = \pm c_A e_{k_1,k'_1} c_{k_1} e_{k_2,k'_2} c_{k_2} \cdots e_{k_t,k'_t} c_{k_t} e_{j_1,j'_1} \cdots e_{j_u,j'_u} c_B \sigma,$$

for some set A, B, $\sigma \in \Sigma$, and j_i, j'_i are all distinct. If there is e_{k_t,k'_t} in $e_{j_1,j'_1} \cdots e_{j_u,j'_u}$, then $w = 0$ by (3),(8). If all j'_i is not k'_t, then we can shift c_{k_t} to the right-hand side of all $e_{p,q}$'s using the relation (6),(7). If there is p such that $j'_p = k'_t$ and $j_p \neq k_t$, then we have

$$w = \pm \cdots e_{k_{t-1},k'_{t-1}} c_{k_{t-1}} c_{j_p} e_{k_t,k'_t} e_{j_1,j'_1} \cdots e_{j_u,j'_u} c_B \sigma.$$

If $j_p = k_{t-1}$, c_{j_p} becomes a constant by relation (5.5). If not, we can shift c_{j_p} to the left-hand side of $e_{k_{t-1},k'_{t-1}}$. In this way, we shift c_{j_p} to the left-side of all $e_{p,q}$'s or c_{j_p} becomes a constant. Repeating this process on $c_{k_1}, \cdots, c_{k_{t-1}}$, we can shift all c_i to the left-hand side or right-hand side of all $e_{p,q}$'s, or obtain $w = 0$.

Combining the relations (5.5), (3),(4),(6)–(8), we finally obtain $w = 0$ or

$$w = \pm c_P e_{p_1,q_1} \cdots e_{p_a,q_a} c_Q \sigma,$$

where $P \subset \{p_1, \cdots, p_a\}$, $Q \subset \{1, \cdots, r+s\} \setminus \{q_1, \ldots, q_a\}$ and $\sigma \in \Sigma$.

Since ψ is surjective, there exists an element $\widetilde{\sigma} \in \Sigma_r \times \Sigma_s$ such that $\psi(\widetilde{\sigma}) = \sigma$. If $\widetilde{\sigma}^{-1}(p_1) > \widetilde{\sigma}^{-1}(p_2)$, we replace σ with $(p_1\ p_2)(q_1\ q_2)\sigma$ using the relation (5). Clearly, $\psi((p_1\ p_2)'(q_1\ q_2)'\widetilde{\sigma}) = (p_1\ p_2)(q_1\ q_2)\sigma$, where $(x\ y)' = \widetilde{s}_x \cdots \widetilde{s}_{y-2} \widetilde{s}_{y-1} \widetilde{s}_{y-2} \cdots \widetilde{s}_x \in \Sigma_r \times \Sigma_s$. In this way, we can find $\sigma \in \Sigma$ satisfying the condition (ii). Therefore, every word in generators of F belongs in the linear span of the elements of X.

The relations (1)–(4) were verified in the proof of [5, Theorem 4.10]. The relation (5.4) implies $e_{r-1,r+2} e_{r,r+1} = e_{r,r+1} e_{r-1,r+2}$.

For the relation (5), we first prove $e_{r,r+1}e_{r-1,r+2}s_{r-1}s_{r+1} = e_{r,r+1}e_{r-1,r+2}$. Note that

$$\begin{aligned}
e_{r,r+1}e_{r-1,r+2}s_{r-1}s_{r+1} &= e_{r,r+1}s_{r-1}s_{r+1}e_{r,r+1} && \text{by (5.1)}\\
&= (e_{r,r+1}s_{r-1}e_{r,r+1})s_{r-1}s_{r+1}e_{r,r+1} && \text{by (5.3)}\\
&= e_{r,r+1}(s_{r+1}s_{r+1})s_{r-1}e_{r,r+1}s_{r-1}s_{r+1}e_{r,r+1} && \text{by (5.1)}\\
&= e_{r,r+1}s_{r+1}e_{r,r+1}s_{r+1}s_{r-1}e_{r,r+1}s_{r-1}s_{r+1} && \text{by (5.4)}\\
&= e_{r,r+1}e_{r-1,r+2} && \text{by (5.3).}
\end{aligned}$$

For $(p,q) \neq (r,r+1), (r-1,r+2)$, we have

$$\begin{aligned}
&e_{p,q}e_{r,r+1}(p\ r)(r+1\ q)\\
&= (r+2\ q)(r-1\ p)e_{r-1,r+2}(r+2\ q)(r-1\ p)e_{r,r+1}(p\ r)(r+1\ q)\\
&= (r+2\ q)(r-1\ p)e_{r-1,r+2}e_{r,r+1}(r+2\ q)(r-1\ p)(p\ r)(r+1\ q)\\
&= (r+2\ q)(r-1\ p)e_{r-1,r+2}e_{r,r+1}(r-1\ r)(r+1\ r+2)(r-1\ p)(r+2\ q)\\
&= (r+2\ q)(r-1\ p)e_{r-1,r+2}e_{r,r+1}(r-1\ p)(r+2\ q)\\
&= e_{p,q}e_{r,r+1}.
\end{aligned}$$

Here, if $p = r-1$ or $q = r+2$, we define $(r-1\ p) = (r+2\ q) = 1$. The general case can be verified in a similar manner.

From the relation (5.7), we obtain

$$\begin{aligned}
c_q e_{p,q} &= c_q(r+1\ q)(r\ p)e_{r,r+1}(r+1\ q)(r\ p) = (r+1\ q)(r\ p)c_{r+1}e_{r,r+1}(r+1\ q)(r\ p)\\
&= (r+1\ q)(r\ p)c_r e_{r,r+1}(r+1\ q)(r\ p) = c_p(r+1\ q)(r\ p)e_{r,r+1}(r+1\ q)(r\ p)\\
&= c_p e_{p,q}.
\end{aligned}$$

The other relations in (6), (7) can be checked similarly.

The relations $e_{p,q}^2 = 0$ and $e_{p,q}c_p e_{p,q} = 0$ follow from (5.1), (5.2) and (5.1), (5.6), (5.8), respectively.

Proof of Step 2:

One can compute the dimension of $\overrightarrow{B}_{r,s}$ by counting the (r,s)-superdiagrams with i horizontal edges in each row and then summing up over all i. Thus we obtain

$$\dim_{\mathbb{C}}(\overrightarrow{B}_{r,s}) = \sum_{i=0}^{\min(r,s)} 2^{r+s}\left(\binom{r}{i}\binom{s}{i}\, i!\right)^2 (r-i)!(s-i)!.$$

Now we count the number of the elements in X. We define a set

$$\begin{aligned} Y = \{(a,&(p_1,q_1),\dots,(p_a,q_a),P,Q,\sigma) \mid \text{(i) } 0 \le a \le \min(r,s),\\ &\text{(ii) } 1 \le p_1 < \cdots < p_a \le r, \text{ and } r+1 \le q_i \le r+s \text{ are all distinct},\\ &\text{(iii) } P \subset \{p_1,\dots,p_a\},\\ &\text{(iv) } Q \subset \{1,\dots,r+s\} \setminus \{q_1,\dots,q_a\},\\ &\text{(v) } \sigma \in \Sigma_r \times \Sigma_s, \text{ and } \sigma^{-1}(p_1) < \cdots < \sigma^{-1}(p_a)\}. \end{aligned}$$

We know $|X| \le |Y|$ and we will count the number of elements of Y. For given $0 \le a \le \min(r,s)$, the number of all distinct subsets of $\{p_1,\dots,p_a\}$ is 2^a and the number of all distinct subsets of $\{1,\dots,r+s\}\setminus\{q_1,\dots,q_a\}$ is 2^{r+s-a}. Observe that the number of distinct $((p_1,q_1),\dots,(p_a,q_a))$'s satisfying the condition (ii) is $\binom{r}{a}\binom{s}{a}a!$. Since $\sigma \in \Sigma_r \times \Sigma_s$ and $\sigma^{-1}(p_1) < \cdots < \sigma^{-1}(p_a)$, the number of σ's satisfying these conditions is $\binom{r}{a}(r-a)!s!$. Therefore

$$|Y| = \sum_{a=0}^{\min(r,s)} 2^{r+s} \left(\binom{r}{a}\binom{s}{a}a!\right)\left(\binom{r}{a}(r-a)!s!\right).$$

From the identity $s! = \binom{s}{a}a!(s-a)!$, we get $|Y| = \dim_{\mathbb{C}}(\overrightarrow{B}_{r,s})$.

Combining Step 1 and Step 2, we obtain $\dim_{\mathbb{C}} F \le |X| \le \dim_{\mathbb{C}}(\overrightarrow{B}_{r,s})$, which completes the proof. □

Remark 5.2.

(1) If $r = 0$ or $s = 0$, then we omit the generator $e_{r,r+1}$ and the corresponding relations.

(2) Since $c_{i+1} = s_i c_i s_i$ for admissible i, the even generators together with c_1, c_{r+1} generate the whole associative superalgebra $\overrightarrow{B}_{r,s}$.

(3) All the relations (5.1)–(5.4) involving $s_i, e_{r,r+1}$ $(i = 1,\dots,r-1,r+1,\dots,r+s-1)$ also appear in the presentation for the walled Brauer algebra; see [6, Corollary 4.5] or [9, Theorem 4.1]. The following relation in [6, Corollary 4.5]

$$s_{r-1}s_{r+1}e_{r,r+1}s_{r+1}s_{r-1}e_{r,r+1} = e_{r,r+1}s_{r-1}s_{r+1}e_{r,r+1}$$

can be derived from the relations (5.1),(5.3) and (5.4).

References

[1] G. Benkart, M. Chakrabarti, T. Halverson, R. Leduc, C. Lee and J. Stroomer, *Tensor product representations of general linear groups and their connections with Brauer algebrs*, J. Algebra **166** (1994), 529–567.

[2] M. Bloss, *G-colored partition algebras as centralizer algebras of wreath products*, J. Algebra **265** (2003), 690–710.

[3] J. Brundan and C. Stroppel, *Gradings on walled Brauer algebras and Khovanov's arc algebra*, Adv. Math. **231** (2012), 709–773.

[4] J. Comes and B. Wilson, *Deligne's category* $\underline{\mathrm{Re}}\mathrm{p}(GL_\delta)$ *and representations of general linear supergroups*, arXiv: 1108.0652.

[5] A. Elduque, *A modified Brauer algebra as centralizer algebra of the unitary group*, Trans. Amer. Math. Soc. **356** (2004), 3963–3983.

[6] T. Halverson, *Characters of the centralizer algebras of mixed tensor representations of* $GL(r,\mathbb{C})$ *and the quantum group* $U_q(\mathfrak{gl}(r,\mathbb{C}))$, Pacific. J. Math. **174** (1996), 359–410.

[7] K. Koike, *On the decomposition of tensor products of the representations of classical groups: by means of universal characters*, Adv. Math. **74** (1989), 57–86.

[8] A. Kleshchev, *Linear and Projective Representations of Symmetric Groups*, Cambridge University Press, Cambridge, 2005.

[9] P. Nikitin, *The centralizer algebra of the diagonal action of the group* $GL_n(\mathbb{C})$ *in a mixed tensor space*, J. Math. Sci. **141** (2007), 1479–1493.

[10] M. Parvathi and M. Kemaraj, *Signed Brauer's algebras*, Comm. Algebra **26** (1998), no.3, 839–855.

[11] M. Parvathi and A.J. Kennedy, *G-vertex colored partition algebras as centralizer algebras of direct products*, Comm. Algebra **32** (2004), 4337–4361.

[12] I. Schur, *Über Eine Klasse Von Matrizen, die Sich Einer Gegeben Matrix Zurodenen Lassen*, Ph.D. thesis (1901). Reprinted in Gesamelte Abhandlungen **1**, 1–70.

[13] I. Schur, *Über die rationalen Darstellungen der allgemeinen linearen Gruppe*, Sitzungsberichte der Königlich Preussischen Adademie der Wissenschaften zu Berlin (1927), 58–75. Reprinted in Gessamelte Abhandlungen **3**, 68–85.

[14] C. L. Shader and D. Moon, *Mixed tensor representations and representations for the general linear Lie superalgebras*, Comm. Algebra **30** (2002), 839–857.

[15] A. Sergeev, *The tensor algebra of the tautological representation as a module over the Lie superalgebras* $\mathfrak{gl}(n,m)$ *and* $Q(n)$, Math. USSR Sbornik **51** (1985), 419–427.

[16] V. Turaev, *Operator invariants of matrices and R-matrices*, Izv. Akad. Nauk SSSR **53** (1989), 1073–1107.

Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Seoul 151-747, Korea

E-mail address: `jhjung@math.snu.ac.kr`

Department of Mathematical Sciences and Research Institute of Mathematics, Seoul National University, Seoul 151-747, Korea

E-mail address: `sjkang@snu.ac.kr`